



## Agromet Advisory Bulletin for Ernakulam District

(Valid from 14.01.2025 to 18.01.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara  
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

**AAS Bulletin No: 02/2025, (4) Monday**

**Dated 13.01.2025**

### A Medium range weather forecast of IMD for next five days

Weather parameters	14.01.2025	15.01.2025	16.01.2025	17.01.2025	18.01.2025
Rainfall (mm)	2	0.1	0.1	0.1	0
Maximum temperature (°C)	32	32	32	32	32
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	76	76	76	80	80
Minimum Relative Humidity (%)	60	60	60	58	58
Wind speed (kmph)	3	3	2	2	2
Wind direction (deg)	320	290	290	320	320
Total cloud cover (octa)	5	4	5	4	3

**Weather summary / Alert:** As per the forecast of IMD, there is a chance of very light rainfall in isolated places of the district. Since dry weather condition is prevails, irrigation of crops can be continued as per requirement. Normal to above normal rainfall is expected in Kerala.

### B. Agrometeorological Advisories for crop protection & management

#### C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

**General Advisory:** - Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Rice Milky stage	Thrips	Flooding the rice field will manage the thrips population to an extent. In severe cases, spray Thiamethoxam 25 WG 2g in 10 litre water.
Vegetables	Mites	2 per cent neem oil + garlic emulsion or commercial neem formulation azadirachtin 0.004 per cent (4 ml per litre of water) or micronized wettable sulphur 0.4 per cent can be applied in the initial stages of infestation. In the case of very severe infestation, the crop can be protected by spraying 0.05% dimethoate or Spiromesifen 22.9% SC 8 ml per 10 litre or Diafenthiuron 50% WP 1.2 g per 10 litre can be sprayed. Rational rotation of pesticides to be followed to avoid resistance development.
Various crops	Caterpillars	Neem seed kernel extract 5% (Grind 50g Neem seed kernel to coarse powder, put in muslin cloth bag, keep immersed in 1 L water overnight. Thereafter, squeeze the cloth bag by dipping in water repeatedly till the water coming out from cloth bag becomes clear); <i>Beauveria bassiana</i> 20g per litre can be sprayed; Chlorantranilipole 18.5 SC 3ml / 10 litre water can be sprayed. If the hairy caterpillars are spreading to homesteads, Cypermethrin 1 ml per litre along with a sticker can be applied.

**Animal Husbandry:** Deworm the animals after dung examination. Vaccinate animals against foot and mouth disease. Care should be taken to avoid ecto parasitic infestation. (CAADECCS, KVASU, Mannuthy).