



Agromet Advisory Bulletin for Ernakulam District

(Valid from 21.01.2023 to 25.01.2023)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 03/2023, (6) Friday

dated 20.01.2023

Weather parameters	21.01.2023	22.01.2023	23.01.2023	24.01.2023	25.01.2023
Rainfall (mm)	0	0	0	0	0.2
Maximum temperature (°C)	31	31	31	31	31
Minimum temperature (°C)	22	22	22	23	23
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	55	55	55	55	55
Wind speed (kmph)	6	6	6	4	3
Wind direction (deg)	250	270	270	250	230
Total cloud cover (octa)	4	4	6	8	8

A. Medium range weather forecast of IMD for next five days

RAINFALL FORECAST FOR NEXT FIVE DAYS				
2023 January 21	2023 January 22	2023 January 23	2023 January 24	2023 January 25
No rain	No rain	No rain	No rain	No rain

Weather summary / Alert: Since dry weather condition is prevailed, irrigation of crops can be continued, when required. Normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulam District (RRS Vyttila)

Since dry weather condition is prevailed, irrigation of crops can be continued, if needed. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Prophylactic measures to control crop pests & diseases can be taken to avoid their spread.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice	Rice bug	There is a chance of appearance of rice bug in paddy. To control, spray 2 ml malathion per one litre of water. Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.
Vegetables	Mites	2 per cent neem oil + garlic emulsion or commercial neem formulation azadirachtin 0.004 per cent (4 ml per litre of water) or micronized wettable sulphur 0.4 per cent can be applied in the initial stages of infestation. In the case of very severe infestation, the crop can be protected by spraying 0.05% dimethoate or Spiromesifen 22.9% SC 8 ml per 10 litre or Diafenthuron 50% WP 1.2 g per 10 litre can be sprayed. Rational rotation of pesticides to be followed to avoid resistance development.
Cauliflower	Management	Cover the curd with 3-4 surrounding leaves to prevent discoloration when the curd is half mature stage (blanching).

Fisheries :-Farmers can culture the following fishes like red snapper, seabass, pearl spot, mullet, pompano, milk fish etc. in brackish water