



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



**ERNAKULAM DISTRICT**

AAS 09/2019(17), Tuesday

26.02.2019

**CURRENT SYNOPTIC SITUATION:** A trough of low at mean sea level lies over Equatorial Indian ocean and adjoining South west Bay of Bengal

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	27.02.2019	28.02.2019	01.03.2019	02.02.2019	03.02.2019
Rainfall (mm)	0	0	0	0	0.6
Maximum temperature (°C )	32	32	32	33	33
Minimum temperature (°C )	25	25	25	24	24
Total cloud cover (octa)	4	4	2	4	6
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	35	35	35	35	35
Wind speed (kmph)	18	16	20	14	18
Wind direction (deg)	270	320	290	270	270

**No rainfall is forecasted for coming five days in the district by India Meteorological Department.**

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Crop stage	Pest/ Disease	Advisories
Coconut	-	Management	Burying of fresh or dried coconut husk around the palm is a desirable practice particularly for moisture retention. The husk can be buried either in linear trenches taken 3 m away from the trunk between rows of palms or in circular trenches taken around the palm at a distance of 2 m from the trunk. The trenches may be of 0.5 m width and depth. The husks are to be placed in layers with concave surface facing upwards and covered with soil. The beneficial effect of husk burial will last for about 5-7 years. Instead of husk, coir pith can be buried @ 25 kg / palm/ year.
Banana	-	Spodoptera	Propping should be followed to reduce the damage due to high wind velocity. Destroy the affected leaves along with the Spodoptera. For severe attack, spray Flubendiamide (Fame) 2 ml/10 L of water.
Vegetables (cowpea)	Fruiting stage	Stunted growth	Due to the increase in atmospheric temperature, the attack of insects such as thrips, mite and white flies etc. are more. As a result of this, there is a chance of stunted growth in vegetables. They can be controlled by applying 10 gms of Verticillium in 1 litre of water or neem based insecticides can be applied.
Animal Husbandry	Animal should not be left to graze under hot sun from morning 10.00 AM to evening 3.00 PM. Ensure sufficient supply of clean drinking water to the animals.(CAADECCS, KVASU, Mannuthy)		