



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Agriculture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



ERNAKULAM DISTRICT

AAS 17/2021, (34)Friday

23.04.2021

CURRENT SYNOPTIC SITUATION: : A cyclonic circulation lies over Marathawada & neighbourhood extending upto 1.5 km above mean sea level. A North-South trough runs from the above cyclonic circulation over Marathawada to south coastal Tamil Nadu across Interior Karnataka extending upto 1.5 km above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	24.04.2021	25.04.2021	26.04.2021	27.04.2021	28.04.2021
Rainfall (mm)	10	10	5	4	5
Maximum temperature(°C)	33	33	33	33	33
Minimum temperature (°C)	26	26	26	26	26
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	66	66	66	66	66
Wind speed (kmph)	10	14	10	14	12
Wind direction (deg)	230	270	270	290	250
Total cloud cover (octa)	7	7	7	5	5

Moderate rainfall is forecasted in isolated places of the district by IMD & KAU. Normal rainfall is expected in Kerala.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: Precaution should be taken during drying of agricultural products and spraying of pesticides.		
Rice	-	Dry sown crop is starting, the land should be prepared well by through ploughing and levelling. Dress the seeds with pseudomonas before sowing. (10 gram of pseudomonas per one kg of paddy seeds.)
Coconut	Management	During the beginning of southwest monsoon season, one-kilogram lime can be applied per one matured palm by opening the coconut basin.
Banana	Sigatoka disease	There is a chance of sigatoka disease in banana. As a precaution, Spray 20 g Pseudomonas per one liter of water.
Ginger, Turmeric	Planting time	While preparing land for ginger and turmeric cultivations apply 2 to 3 kg lime per cent. As a prophylactic measure against foot rot disease, the ginger and turmeric rhizome should be soaked in the 0.3 percent mancozeb or pseudomonas solution (3 gram per one litter of water) for 30 minutes. Moreover 90 kg FYM mixed with one kg Trichoderma and kept under shade for two weeks with sufficient moisture and can be applied.
Animal Husbandry		Assure increased ventilation in the shed.