





## Agromet Advisory Bulletin for Ernakulam District (Valid from 11.6.2022 to 15.6.2022)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 23/2022, (46) Friday dated 10.6.2022

## A. Medium range weather forecast of IMD for next five days

Weather parameters	11.6.2022	12.6.2022	13.6.2022	14.6.2022	15.6.2022
Rainfall (mm)	10	10	12	4	2
Maximum temperature (°C)	31	30	30	30	30
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	89	85	89	86	89
Minimum Relative Humidity (%)	57	57	58	57	62
Wind speed (kmph)	3	3	3	3	3
Wind direction (deg)	230	270	230	290	290
Total cloud cover (octa)	7	8	8	4	6

RAINFALL FORECAST FOR NEXT FIVE DAYS						
2022 June 11	2022 June 12	2022 June 13	2022 June 14	2022 June 15		
Light rain	Light rain	Light rain	Light rain	Very light rain		

**Weather summary / Alert:** Light rainfall is expected in the district and moderate rainfall is expected in isolated places. Below normal rainfall is expected in Kerala.

## B. Agrometeorological Advisories for crop protection & management

## C. General and Agrometeorological Advisories for (RRS Vyttila)

Proper drainage facilities have to be ensured. Special crop management is required for agricultural crops during monsoon season. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rain.

iuni.							
Crop	Pest/ Disease		Crop/ Animal Advisory				
Paddy (Pokkali)	Management	After checking the salinity, preparations for sowing can be started					
Paddy (Virippu)	Management	Preparations for sowing can be done in the case of direct sowing or Sowing in nursery for transplantation.					
Coconut	Bud rot	the crov Bordeau small pe	e general condition of the palm by manuring and management. Clean wn and remove rotten portions. Spray crowns and leaves with 1 % ax mixture as a prophylactic measure. For already infected palms, erforated sachets containing 5 g Mancozeb may be tied to the top portion and 3 or 4 inner most leaf axils.				
Pepper	Foot rot	As a prophylactic measure, add 2.5 kg Trichoderma enriched organic man to each vine.  As a precautionary measure, spray 1% Bordeaux mixture					
Nutmeg	Leaf fall						
Fisheries			hare ponds for fish farming. Drain the pond and apply lime. Apply teake in ponds where draining is difficult to eradicate weed fishes.				