



Agromet Advisory Bulletin for Ernakulam District

(Valid from 12.06.2024 to 16.06.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 24/2024, (47) Tuesday

Dated 11.06.2024

A Medium range weather forecast of IMD for next five days

Weather parameters	12.06. 2024	13.06.2024	14.06. 2024	15.06. 2024	16.06.2024
Rainfall (mm)	6	5	3	1	1
Maximum temperature (°C)	31	32	32	32	32
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	93	93	93	93	93
Minimum Relative Humidity (%)	82	82	82	82	82
Wind speed (kmph)	8	6	2	3	4
Wind direction (deg)	320	320	340	270	320
Total cloud cover (octa)	8	8	8	8	8

Weather summary / Alert: Thunderstorm with lightning is very likely to occur at one or two places in Kerala on 11 th & 12 th June 2024.

SMS: - As per the forecast of IMD, there is a chance of Moderate rainfall in isolated places of the district. Normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Special crop management is required for agricultural crops before monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports for the bananas. Make the vegetable pandals stronger. Avoid standing in open areas to protect farmers and livestock from lightning and thunder.

Crop	-	Crop/ Animal Advisory
Paddy (Pokkali)	Management	The sprouted seeds to be sown on the top of the already prepared mounds and the seeds to be plastered with mud.
Paddy (Virippu)	Management	Preparations for sowing can be done in the case of direct sowing or sowing in nursery for transplantation.
Banana	Sigatoka leaf spot	Remove and destroy severely infected and completely dried leaves. Improve drainage facilities. Spray 1 % Bordeaux mixture or Mancozeb 75 WP 3g per litre water along with stickers 4-5 times at 2-3 weeks interval as a prophylactic measure. In diseased field spray Hexaconazole 5EC ml per litre water.
Pepper	Foot rot	As a prophylactic measure, add 2.5 kg Trichoderma enriched organic manure to each vine.
Vegetables	Leaf eating caterpillars	Collection and destruction of pest infected parts along with caterpillars, spray neem oil emulsion or neem seed kernel extract or Beauveria 20g / litre of water in the initial stages of infestation itself.

Animal Husbandry: - Deworm the animals after dung examination. Take precautions to avoid ecto parasitic infestation. Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospirosis..