





## Agromet Advisory Bulletin for Ernakulam District (Valid from 14.06.2023 to 18.06.2023)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

#### AAS Bulletin No: 24/2023, (47) Tuesday

Dated 13.06.2023

## A. Medium range weather forecast of IMD for next five days

Weather parameters	14.06.2023	15.06.2023	16.06.2023	17.06.2023	18.06.2023
Rainfall (mm)	4	13	5	14	12
Maximum temperature (°C)	32	31	32	32	32
Minimum temperature (°C)	26	24	26	25	26
Maximum Relative Humidity (%)	90	90	88	92	88
Minimum Relative Humidity (%)	64	68	62	70	70
Wind speed (kmph)	5	3	3	5	4
Wind direction (deg)	270	290	270	290	290
Total cloud cover (octa)	7	8	6	8	8

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in the district. Normal rainfall is expected in Kerala.

# B. Agrometeorological Advisories for crop protection & management

### C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rain.

The state of the s	not of appared during runn.				
Crop	Pest/ Disease	Crop/ Animal Advisory			
Paddy	Managamant	After checking the salinity, preparations for sowing can be			
(Pokkali)	Management	done			
Paddy	Managamant	Preparations for sowing can be done in the case of direct			
(Virippu)	Management	sowing or sowing in nursery for transplantation.			
Banana		Remove and destroy severely infected and completely dried			
		leaves. Improve drainage facilities. Spray 1 % Bordeaux			
	Sigatoka leaf	mixture or Mancozeb 75 WP 3g per litre water along with			
	spot	stickers 4-5 times at 2-3 weeks interval as a prophylactic			
	-	measure. In diseased field spray Hexaconazole 5EC ml per			
		litre water.			
Pepper	Foot rot	As a prophylactic measure, add 2.5 kg Trichoderma enriched			
		organic manure to each vine.			
Vegetables		Collection and destruction of pest infected parts along with			
	Leaf eating	caterpillars, spray neem oil emulsion or neem seed kernel			
	caterpillars	extract or Beauveria 20g / litre of water in the initial stages of			
	_	infestation itself.			