



Agromet Advisory Bulletin for Ernakulam District

(Valid from 24.06.2023 to 28.06.2023)

*Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi*

AAS Bulletin No: 25/2023, (50) Friday

Dated 23.06.2023

A. Medium range weather forecast of IMD for next five days

Weather parameters	24.06.2023	25.06.2023	26.06.2023	27.06.2023	28.06.2023
Rainfall (mm)	2	3	4	8	25
Maximum temperature (°C)	32	31	32	32	32
Minimum temperature (°C)	24	26	26	26	26
Maximum Relative Humidity (%)	90	87	87	89	95
Minimum Relative Humidity (%)	75	65	63	62	69
Wind speed (kmph)	5	8	5	4	5
Wind direction (deg)	340	320	290	270	290
Total cloud cover (octa)	8	8	8	8	8

Weather summary / Alert: As per the forecast of IMD, there is a chance of Moderate rainfall in the district. Above normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulam District (RRS Vyttila)

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rain.

Crop	Pest/ Disease	Crop/ Animal Advisory
Paddy (Pokkali)	Management	After checking the salinity, preparations for sowing can be done
Paddy (Virippu)	Management	Preparations for sowing can be done in the case of direct sowing or sowing in nursery for transplantation.
Coconut	Management	Ensure drainage, Apply 25-50 Kg organic manure after preparing basins with 1.8 m radius and 25 cm depth, coconut husk burial can be practiced to harvest maximum rain water, seeds of green manure crops can be sown in the basins.
Banana	Sigatoka leaf spot	Remove and destroy severely infected and completely dried leaves. Improve drainage facilities. Spray 1 % Bordeaux mixture or Mancozeb 75 WP 3g per litre water along with stickers 4-5 times at 2-3 weeks interval as a prophylactic measure. In diseased field spray Hexaconazole 5EC ml per litre water.
Pepper	Planting	Two rooted cuttings each can be planted in the pits prepared in advance