

## Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in





(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)

## ERNAKULAM DISTRICT

AAS 27/2020,(54) Friday 03.07.2020

CURRENT SYNOPTIC SITUATION: The cyclonic circulation over Westcentral & adjoining Northwest Arabian Sea extending upto 5.8 km above mean sea level persists. The off-shore trough from north Maharashtra coast to Karnataka coast runs from north Maharashtra coast to north Kerala coast. A cyclonic circulation lies over Westcentral Bay of Bengal off Andhra Pradesh coast at 3.1 km above mean sea level.

## WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	04.07.2020	05.07.2020	06.07.2020	07.07.2020	08.07.2020
Rainfall (mm)	20	15	8	8	6
Maximum temperature(°C)	31	31	31	31	31
Minimum temperature (°C)	24	24	24	24	24
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	6	8	6	8	6
Wind direction (deg)	290	290	320	320	320
Total cloud cover (octa)	8	8	8	8	7

As per the forecast of IMD & KAU, there is a chance of decreasing the intensity of rainfall.

AGRO-METEOROLOGICAL ADVISORIES:				
Crop	Pest/ Disease	Advisories		
General Advisories: Proper drainage facilities should be ensured to protect the agricultural				
crops from pest & diseases during rainy season.				
Paddy	Seedling stage	During rainy season, the micronutrient deficiency of paddy can		
		be managed by spraying 10 gram of paddy sampoorna in one		
		litre of water.		
Coconut	Management	During the begining of southwest monsoon season, one-		
		kilogram lime can be applied per one matured palm by opening		
		the coconut basin.		
Banana	Pseudo stem	There is a chance of pseudo stem weevil attack in banana. As		
	weevil	a precaution, apply 50 g neem cake powder in leaf axils.		
Vegetables	-	During rainy season, the micronutrient deficiency of vegetables		
		can be managed by spraying 5 gram of vegetable sampoorna in		
		one litre of water.		
	Keep the cattle shed and premises clean to avoid mosquito breeding. Avoid			
Animal Husbandry	waterlogging near the cattle shed. The cowdung pit can be covered to protect from			
	direct falling of rain.			
	direct failing of faili.			