


## Agromet Advisory Service bulletin 28/17, Tuesday dt 11.07.2017

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



## **ERNAKULAM**

**CURRENT SYNOPTIC SITUATION:** An off shore trough runs from South Maharashtra coast to North Kerala coast.

## NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 11.07.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 12/07	DAY-2 13/07	DAY-3 14/07	DAY-4 15/07	DAY-5 16/07
DISTRICT : ERNAKULAM					
ainfall (mm)	3	3	11	10	4
lax Temperature ( deg C)	32	31	32	32	32
in Temperature ( deg C)	24	24	24	24	24
otal cloud cover (octa)	5	5	7	6	5
ax Relative Humidity (%)	90	90	90	90	90
in Relative Humidity (%)	80	80	80	80	80
ind speed (kmph)	014	014	014	014	014
ind direction (deg)	290	290	290	270	270

## **Agro Meteorological Advisories:**

Crop	Stage of the crop / Pest/Disease	Advisories
		There is a chance of rice case worm in paddy field. To
Paddy	Seedling stage/Rice	control this 25 kg sawdust or rice husk ash mixed with 1
(Jyothi, Kanchana)	case worm	litre kerosene can be spread after draining the field for 1
		acre.
Banana		Due to the pre monsoon showers, there is chance of
		Sigatoka disease in banana. As a precaution, Spray
	Sigatoka disease	Pseudomonas 20g/L In case of severe attack spray 1 g
		Bavistin by mixing it with gum in 1 litre of water on lower
		surface of leaf.
Vegetables	Aphids	Apply tobacco decoction or 2% neem oil-garlic-soap emulsion for control of aphids in cowpea.
<u> </u>		As part of the mosquito control measures, the direct rain
		falling on the cowdung pit should be avoided.
Animal Husbandry		
	-	Thank you for Information: Centre for animal adaptation
		environment and climate change studies, KVASU,
		Mannuthy.