



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



ERNAKULAM DISTRICT

AAS 30/2020,(60)Friday

24.07.2020

CURRENT SYNOPTIC SITUATION:A cyclonic circulation lies over Eastcentral Arabian sea off North Kerala - Karnataka coast extending upto 3.6 km above mean sea level.

HEAVY RAINFALL WARNING:

DAY 1: Heavy rainfall (7- 11 cm in 24 hours) is very likely to occur at one or two places in the districts of Alappuzha, Eranakulam, Idukki , Malappuram, Kozhikode, Kanuur and Kasaragod.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	25.07.2020	26.07.2020	27.07.2020	28.07.2020	29.07.2020
Rainfall (mm)	22	11	12	15	16
Maximum temperature(°C)	31	31	31	31	31
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	95	95	95	95	95
Minimum Relative Humidity (%)	73	73	73	73	73
Wind speed (kmph)	18	22	14	20	18
Wind direction (deg)	230	230	320	320	320
Total cloud cover (octa)	8	8	8	8	8

As per the forecast of IMD & KAU, there is a chance of moderate rainfall in the district and a chance of heavy rainfall in isolated places.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: Proper drainage facilities should be ensured to protect the agricultural crops from pest & diseases during rainy season.		
Rice (Virippu) Tillering stage	Bacterial Leaf Blight (BLB)	As a prophylactic measure to manage Bacterial Leaf Blight during rainy season, spray a mixture of cow dung slurry (20g cow dung per one litre of water) and pseudomonas solution (20g per one litre of water).
Banana	Rhizome rot	Due to the increase in rainfall, there is a chance of Rhizome rot in banana. As a precaution, drench the basin with three litre of pseudomonas solution (20g per one litre of water).
Pepper	Foot rot	As a prophylactic measure to manage foot rot during rainy season, add 2 kg Trichoderma to a mixture of 90 kg FYM &10 kg neem cake and kept for two weeks with sufficient moisture. Add 2.5 kg of this mixture to each matured pepper vine to control foot rot disease.
Cucurbitaceous Vegetables	Downy mildew	Due to high relative humidity, there is a chance of downy mildew attack in cucurbitaceous vegetables. As a prophylactic measure, spray pseudomonas solution (20 g per one litre) under the leaves.
Animal Husbandry	Don't allow the animals to water logged areas to avoid foot lesions.	