

Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



ERNAKULAM DISTRICT

AAS 31/2020,(62) Thursday

30.07.2020

CURRENT SYNOPTIC SITUATION: A cyclonic circulation lies over South coastal Andhra Pradesh and adjoining North Tamilnadu between 3.1 & 5.8 km above mean sea level tilting southwestwards with height.

HEAVY RAINFALL WARNING:

DAY 1: Heavy rainfall (7-11 cm in 24 hours) is very likely to occur at one or two places in the districts of Kollam, Alappuzha, Kottayam, Eranakulam, Thrissur & Palakkad.

DAY 2: Very Heavy rainfall (12- 21 cm in 24 hours) is very likely to occur at one or two places in the district of Alappuzha Kottayam, Eranakulam, & Idukki.

DAY 4: Heavy rainfall (7-11 cm in 24 hours) is very likely to occur at one or two places in the districts of Alappuzha, Eranakulam, . Idukki Thrissur, Malappuram, Kozhikode, Kanuur & Kasaragod.

DAY 5: Heavy rainfall (7-11 cm in 24 hours) is very likely to occur at one or two places in the districts of of Alappuzha, Kottayam, Eranakulam, . Idukki Malappuram, Wayanad, Kozhikode, Kanuur & Kasaragod.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	31.07.2020	01.08.2020	02.08.2020	03.08.2020	04.08.2020
Rainfall (mm)	32	45	25	23	20
Maximum temperature(°C)	28	30	30	30	30
Minimum temperature (°C)	23	24	24	24	24
Maximum Relative Humidity (%)	98	98	98	98	98
Minimum Relative Humidity (%)	78	78	78	78	78
Wind speed (kmph)	18	20	18	20	18
Wind direction (deg)	270	270	290	290	320
Total cloud cover (octa)	8	8	8	8	8
		•		•	

As per the forecast of IMD & KAU, there is a chance of moderate rainfall in the district.

AGRO-METEOROLOGICAL ADVISORIES:

AURO-MIETEUROE	AGRO-METEOROLOGICAL ADVISORIES:				
Crop	Pest/ Disease	Advisories			
General Advisories: Proper drainage facilities should be ensured to protect the agricultural crops from pest &					
diseases during rainy season.					
Rice	Bacterial Leaf	As a prophylactic measure to manage Bacterial Leaf Blight during rainy			
(Virippu)	Blight (BLB)	season, spray a mixture of cow dung slurry (20g cow dung per one litre of			
Tillering stage	Blight (BLB)	water) and pseudomonas solution (20g per one litre of water).			
Banana	Panama wilt	There is a chance of panama wilt in banana varieties like poovan, kadali etc.			
		Drench the soil with one gram of carbendazim in one litre of water.			
Vegetables	Anthracnose	During rainy season, there is a chance of Anthracnose disease in Cowpea. As			
(Cowpea)		a prophylactic measure, apply 1% Bordeaux mixture or spray copper			
	disease	oxychloride 3 gram per one litre of water.			
Animal	Animal Deworming of animals should be done only after dung sample examination. In calves first				
Husbandry	dose of medicine should be given within 15 days of age.				