



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Agriculture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



ERNAKULAM DISTRICT

AAS 32/2021, (63)Tuesday

03.08.2021

CURRENT SYNOPTIC SITUATION: A Well Marked Low Pressure Area lies over northwest Madhya Pradesh & neighborhood with associated cyclonic circulation extending up to 7.6 km above mean sea level. A cyclonic circulation lies over north Bay of Bengal & neighborhood between 1.5 & 5.8 km above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	04.08.2021	05.08.2021	06.08.2021	07.08.2021	08.08.2021
Rainfall (mm)	7	11	10	10	11
Maximum temperature(°C)	30	30	30	30	31
Minimum temperature (°C)	24	24	24	24	25
Maximum Relative Humidity (%)	92	92	92	92	92
Minimum Relative Humidity (%)	72	72	72	72	72
Wind speed (kmph)	14	20	20	20	16
Wind direction (deg)	320	320	320	320	320
Total cloud cover (octa)	7	7	8	8	8

As per the forecast of IMD & KAU, there is a chance of light rainfall in the district. Below normal rainfall is expected in Kerala.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing.		
Paddy (virippu) Tillering stage	Gall midge	There is a chance of Gall midge attack in rice. To control this apply 6 kilogram of Carbosulfan per acre.
Banana	Rhizome rot	Due to the increase in rainfall, there is a chance of Rhizome rot in banana. As a precaution, pour 3 litre of pseudomonas solution (20 gm per 1 litre) at the basin.
Pepper	Foot rot	As a prophylactic measure to manage foot rot during rainy season, add 2 kg Trichoderma to a mixture of 90 kg FYM & 10 kg neem cake and kept for two weeks with sufficient moisture. Add 2.5 kg of this mixture to the base of each matured pepper vine to control foot rot disease.
Nutmeg	Leaf fall	As a precautionary measure to manage leaf fall during rainy season, spray 1% bordeaux mixture.
Ginger	Stem borer	There is a chance of stem borer in ginger. Spray 20 g <i>Beauveria</i> per one litre of water to control stem borer.
Turmeric	Leaf Blotch	During rainy season there is a chance of Leaf blotch in Turmeric. To control, spray 2 ml. Contaf (Hexaconazole) in one litre of water.
Vegetables (Bindi)	Leaf spot	There is a chance of leaf spot attack in bindi. To control leaf spot, spray three gram mancozeb per litre of water.
Animal Husbandry	Sun dry the cattle feed frequently to avoid fungal growth. Give foot and mouth disease (FMD) vaccination to the animal (caadeccs, kvasu, Mannuthy).	

(Information given by RRS Vyttila)