



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**  
**Email: cohagmet@kau.in&kauagmet@yahoo.co.in**



**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)

**ERNAKULAM DISTRICT**

**AAS 36/2019(72), Friday**

**06.09.2019**

**CURRENT SYNOPTIC SITUATION:** A Low pressure area lies over northCoastal Odisha & adjoining Gangetic West Bengal with associated cyclonic circulation extending upto7.6km above mean sea level tilting southwestwards with height.An offshore trough at mean sea level runs from Gujarat coast to North Kerala coast.An East- west shear zone lies roughly along Lat 21° N between 3.1 and 7.6 km above mean sea level tilting southwards with height .

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

<b>Weather parameters</b>	<b>07.09.2019</b>	<b>08.09.2019</b>	<b>09.09.2019</b>	<b>10.09.2019</b>	<b>11.09.2019</b>
Rainfall (mm)	35	30	25	20	15
Maximum temperature(°C )	31	31	31	31	31
Minimum temperature (°C )	24	24	24	24	24
Maximum Relative Humidity (%)	96	96	96	96	96
Minimum Relative Humidity (%)	78	78	78	78	78
Wind speed (kmph)	18	16	18	20	12
Wind direction (deg)	290	290	290	320	290
Total cloud cover (octa)	8	8	8	7	7

**According to the forecast of IMD & KAU, as per the current situation, moderate rainfall is expected in coming days in the district.**

**AGRO-METEOROLOGICAL ADVISORIES:**

<b>Crop</b>	<b>Pest/ Disease</b>	<b>Advisories</b>
<b>General Advisories:</b> Proper drainage facilities should be ensured to protect the agricultural crops from crop damage as well as pest & diseases during rainy season.		
Rice( <i>Virippu</i> ) Milky stage	Rice bug	There is a chance of appearance of rice bug in paddy. To control them, spray 2 ml malathion per one litre of water. Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.
Coconut	Bud rot disease	There is a chance of bud rot in rainy season. Keep three numbers each of perforated Mancozeb sachets (5 gm per packet) around spindle leaves as a prophylactic measure against bud rot.
Nutmeg	Fruit rot & leaf fall	During rainy season, there is a chance of Fruit rot and leaf fall in Nutmeg. As a prophylactic measure, apply <i>Trichoderma</i> mixture. Add 2 kg <i>Trichoderma</i> to a mixture of 90 kg FYM and 10 kg Neemcake and kept for two weeks with sufficient moisture. Basal application of 2.5 kg of this mixture will be followed by foliar spray with pseudomonas (20g/litre) or 1% Bordeaux mixture.
Ginger, Turmeric	Rhizome rot	During rainy season there is a chance of rhizome rot. To control this, dig out the affected plants and drench the soil with copper oxychloride 2g per one litre of water.
Vegetables (cowpea)	Anthracoese disease	During rainy season, there is a chance of Anthracnose disease in vegetables. As a prophylactic measure, apply 1% Bordeaux mixture.
Animal Husbandry		Avoid water logging in the premises of cattle shed. Take measures for avoid housefly and mosquito breeding. (CAADECCS, KVASU Mannuthy)