



Agromet Advisory Bulletin for Ernakulam District

(Valid from 11.09.2024 to 15.09.2024)

*Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi*

AAS Bulletin No: 37/2024, (73) Tuesday

Dated 10.09.2024

A Medium range weather forecast of IMD for next five days

Weather parameters	11.09.2024	12.09.2024	13.09.2024	14.09.2024	15.09.2024
Rainfall (mm)	3	2	3	4	3
Maximum temperature (°C)	31	32	32	32	32
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	90	90	88	92	92
Minimum Relative Humidity (%)	80	80	76	86	86
Wind speed (kmph)	4	5	4	4	5
Wind direction (deg)	270	270	290	290	290
Total cloud cover (octa)	8	8	5	8	8

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in isolated places of the district. Normal to above normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports to banana crops. Make the vegetable pandals stronger. To protect farmers and livestock from lightning and thunder avoid standing in open areas.

Crop	-	Crop/ Animal Advisory
Paddy (Pokkali)	Stem borer	Release of egg parasitoids <i>Trichogramma japonicum</i> (5cc / Ha) at weekly intervals from 15-30 days after transplantation immediately after noticing stem borer activity.
Paddy (Virippu)	Rice bug	Fish amino acid @ 20 ml /L water or Azadirachtin (3000 ppm) 10 ml / L water can be sprayed. Farmers those who cultivates for second crop can initiate preparations for nursey.
Coconut	Bud rot	Ensure drainage. Prophylatic management practices against bud rot to be taken
Ginger	Soft rot & stem borer	Spray <i>Beauveria bassiana</i> and <i>Pseudomonas</i> @ 20g in a litre of water in alternate weeks

Animal Husbandry: - Cattle feed should be stored on a wooden board or iron board. If moisture is in the cattle feed it may cause deadly fungal growth. Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospirosis. (CAAECCS, KVASU, Mannuthy).