



Agromet Advisory Bulletin for Ernakulam District

(Valid from 12.10.2022 to 16.10.2022)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 41/2022, (81) Tuesday **dated 11.10.2022**

A. Medium range weather forecast of IMD for next five days

Weather parameters	12.10.2022	13.10.2022	14.10.2022	15.10.2022	16.10.2022
Rainfall (mm)	2	0.4	1	0.5	4
Maximum temperature (°C)	31	31	31	31	31
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	83	83	83	83	83
Minimum Relative Humidity (%)	73	73	73	73	73
Wind speed (kmph)	4	3	3	2	2
Wind direction (deg)	250	270	270	250	200
Total cloud cover (octa)	7	5	8	7	8

RAINFALL FORECAST FOR NEXT FIVE DAYS				
2022 October 12	2022 October 13	2022 October 14	2022 October 15	2022 October 16
Very light rain	Very light rain	Very light rain	Very light rain	Light rain

Weather summary / Alert: Very light rainfall is expected in the district. Normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District

Prophylactic measures to control crop pest & diseases can be taken to avoid their spread.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice (Mundakan)	Land preparation	Land preparations for mundakan paddy can be started in kole paddy field, apply lime during nursery preparation. Seed treatment with <i>pseudomonas</i> can be done before sowing. (10 gram of <i>pseudomonas</i> for one kg of paddy seeds.)
Banana	Planting time	The rhizomes are to be smeared with cowdung solution & ash then they are dried in the sun for about 3 to 4 days before planting. Lime can be applied at 250 gram per plant at the time of planting.
Turmeric	Leaf Blotch (<i>Taphrina maculans</i>)	As a prophylactic measure, spray 2ml hexaconazole or 1 ml Propiconazole or 2.5 gram copper oxychloride or 2 gram SAAF dissolved in one litre of water. Spraying can be continued at fortnightly intervals depending upon the intensity of the disease.

Animal Husbandry- Precautions to be taken to avoid leptospirosis and ectoparasitic infestations. Care should be taken to avoid fungal growth in cattle feed. (CAAECCS, KVASU, Mannuthy).