





#### Agromet Advisory Bulletin for Ernakulam District (Valid from 25.10.2023 to 29.10.2023)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

#### AAS Bulletin No: 43/2023, (85) Tuesday

Dated 24.10 .2023

## A. Medium range weather forecast of IMD for next five days

Weather parameters	25.10.2023	26.10.2023	27.10.2023	28.10.2023	29.10.2023
Rainfall (mm)	8	7	3	4	6
Maximum temperature (°C)	31	31	31	31	31
Minimum temperature (°C)	24	24	24	24	24
Maximum Relative Humidity (%)	95	95	95	95	95
Minimum Relative Humidity (%)	65	65	65	65	65
Wind speed (kmph)	4	4	4	4	4
Wind direction (deg)	250	50	270	250	250
Total cloud cover (octa)	8	8	7	8	8

Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in the district and moderate rainfall in isolated places. Normal rainfall is expected in Kerala.

### **B.** Agrometeorological Advisories for crop protection & management

# C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila) Prophylactic measures to control crop pest & diseases can be taken to avoid their spread. Proper mulching can be followed to conserve soil moisture.

Paddy (Pokkali)ManagementMeasures for dewatering of excess water to be done. Reflective ribbons can be tied up above the crop canopy to ward off purple moorhens' damage.Paddy (Mundakan)ManagementContinue land preparation, basal fertilizer application, transplantation etc.Pool season vegetablesPlanting and managementContinue land preparation etc.Cool season vegetablesPlanting and managementThe season for cool season vegetables is from October to February. Well drained loose soil is preferred. Apply 2 Kg lime at the time of land preparation. Incorporate 100 Kg <i>Trichoderma</i> enriched farm yard manure / compost two weeks after the application of lime. Seedlings can be transplanted by the last of October or first week of November.Fisheries: - As there are chances of PH fluctuations in pond water, apply lime on the bunds after testing the pond water. Rice research Station, Vyttila (Kerala Agricultural University) is	Сгор	Pest/ Disease	Crop/ Animal Advisory		
(Mundakan)Managementtransplantation etc.(Mundakan)The season for cool season vegetables is from October to February. Well drained loose soil is preferred. Apply 2 Kg lime at the time of land preparation. Incorporate 100 Kg <i>Trichoderma</i> enriched farm yard manure / compost two weeks 	2	Management	ribbons can be tied up above the crop canopy to ward off		
Cool season vegetablesPlanting and managementFebruary. Well drained loose soil is preferred. Apply 2 Kg lime at the time of land preparation. Incorporate 100 Kg <i>Trichoderma</i> enriched farm yard manure / compost two weeks after the application of lime. Seedlings can be transplanted by the last of October or first week of November.Fisheries: - As there are chances of PH fluctuations in pond water, apply lime on the bunds	2	Management			
		U	February. Well drained loose soil is preferred. Apply 2 Kg lime at the time of land preparation. Incorporate 100 Kg <i>Trichoderma</i> enriched farm yard manure / compost two weeks after the application of lime. Seedlings can be transplanted by		
$\pm a \pi e_1 = e_1 \pi e_2 \pi e_1 + e_2 \pi e_1 + e_1 + e_2 \pi e_2 \pi e_1 + e_1 + e_2 \pi e_1 + e_1 + e_2 \pi e_2 \pi e_1 + e_2 \pi e_2 \pi e_2 \pi e_1 + e_2 \pi e_2 $					

having facilities to test water quality parameters of fish culture ponds.