

Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in







ERNAKULAM DISTRICT

AAS 47/2018(92), Tuesday

21.11.2018

CURRENT SYNOPTIC SITUATION: The well marked low pressure area over south west Bay of Bengal off Tamil Nadu coast now lies over southwest Bay of Bengal and adjoining coastal Tamil Nadu with associated upper air cyclonic circulation extending upto 5.8 km above mean sea level. A trough from this cyclonic circulation to Maldives area across Interior Tamil Nadu and Kerala between 2.1 and 3.1 km above mean sea level. The well marked low pressure area over south west Arabian sea with associated upper air cyclonic circulation extending upto 4.5 km above mean sea level persists.

HEAVY RAINFALL WARNING:

DAY 1& 2: Heavy rainfall is very likely to occur at one or two places in Kerala.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	22.11.2018	23.11.2018	24.11.2018	25.11.2018	26.11.2018
Rainfall (mm)	8	17	18	20	10
Maximum temperature (°C)	31.1	31.1	31.1	31.1	31.1
Minimum temperature (°C)	23.1	23.1	23.1	23.1	23.1
Total cloud cover (octa)	5	7	7	7	5
Maximum Relative Humidity (%)	95	95	95	95	95
Minimum Relative Humidity (%)	75	75	75	75	75
Wind speed (kmph)	6	6	6	6	6
Wind direction (deg)	140	50	130	230	230

As per IMD forecast, moderate rainfall is expected for coming week in the district.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Crop stage	Pest/ Disease	Advisories	
Paddy	Seedling stage	Rice armyworm	 There is a chance of rice armyworm in the paddy field. To control, Complete flooding of field for 24 hours wherever possible. Spraying of chlorantraniliprole @ 3 ml per 10 litre of water in the field, bunds and weeds. In the fields where transplanting is not yet undertaken, application of chlorantraniliprole granules @ 4 kg per one acre along with basal dose of fertilizers. Keep the field bunds and surroundings weed free. 	
Vegetables (cowpea)	-	Aphids	There is a chance of aphids in cowpea. Apply 2% neemoil emulsion or apply 20 g verticillium lecanii per one litre of water in 10 days intervals.	
			If aphid attack is severe, apply 3 ml Imidacloprid or 2 g Thiamethoxam per 10 litre of water.	
Banana	-	Spodoptera	Destroy the affected leafs along with the <i>Spodoptera</i> . If <i>Spodoptera</i> attack is severe, spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml per 10 litre of water.	
Animal Husbandry	Mix 30g sodium bicarbonate and 1-teaspoon yeast in the feed daily to avoid digestive disorders associated with excessive acidity. (CAADECCS, KVASU, Mannuthy)			