



Agromet Advisory Bulletin for Ernakulam District

(Valid from 07.12.2024 to 11.12.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 49/2024, (98) Friday

Dated 06.12.2024

A Medium range weather forecast of IMD for next five days

Weather parameters	07.12.2024	08.12.2024	09.12.2024	10.12.2024	11.12.2024
Rainfall (mm)	1	2	0.5	0.4	1
Maximum temperature (°C)	31	31	31	31	31
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	86	86	86	86	86
Minimum Relative Humidity (%)	76	76	76	76	76
Wind speed (kmph)	2	2	3	3	3
Wind direction (deg)	270	290	290	270	270
Total cloud cover (octa)	4	3	4	4	5

Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in the district. Normal to above normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Provide sufficient drainage facilities in agricultural lands. Spraying should be done only on dry/clear weather conditions. Do not keep the harvested produces like rice, rubber and copra at open places.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Rice (General)	Bacterial Leaf blight	Dissolve 20 g cow dung in one litre of water and spray supernatant liquid by mixing with 20 g <i>Pseudomonas</i> in one litre of water.
Coconut	Rhinoceros beetle / Red Palm weevil	Field sanitation; top most three leaf axils may be filled with 25g neem cake and 200g sand mixture or naphthalene balls and sand as a prophylactic measure and repeat this procedure 45 days interval; Crown cleaning followed by the application of 20g cartap hydrochloride 4G/ Fipronil 0.3G / Carbosulphan 6G mixed with 200g sand on top three leaf axils; mist net balls can be inserted around leaf spindles.
Banana	Leaf spot	20g / L <i>Pseudomonas</i> can be sprayed in initial stages, rhizome treatment in 50g <i>Pseudomonas</i> per litre water for 30 minutes; 1% Bordeaux mixture can be sprayed 3 to 4 times in 3 to 4 weeks interval.
Cool season vegetables	Management	Fertilizer application

Animal Husbandry: - Vaccinate animals against foot and mouth disease and hemorrhagic septicemia. If needed give medicine against worm infestation after dung examination. (CAAECCS, KVASU, Mannuthy).