



## Agromet Advisory Bulletin for Ernakulam District

(Valid from 28.12.2022 to 01.01.2023)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara  
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 52/2022, (103) Tuesday

dated 27.12.2022

### A. Medium range weather forecast of IMD for next five days

Weather parameters	28.12.2022	29.12.2022	30.12.2022	31.12.2022	01.01.2023
Rainfall (mm)	4	0.5	0	0	0
Maximum temperature (°C)	32	32	32	32	31
Minimum temperature (°C)	25	25	25	25	23
Maximum Relative Humidity (%)	89	89	89	89	89
Minimum Relative Humidity (%)	67	67	67	67	67
Wind speed (kmph)	8	8	6	6	6
Wind direction (deg)	230	270	270	250	250
Total cloud cover (octa)	8	8	5	6	6

### RAINFALL FORECAST FOR NEXT FIVE DAYS

2022 December 28	2022 December 29	2022 December 30	2022 December 31	2023 January 01
Light rain	Very light rain	No rain	No rain	No rain

Weather summary / Alert: Light rainfall is expected in the district. Above normal rainfall is expected in Kerala.

### B. Agrometeorological Advisories for crop protection & management

#### C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Prophylactic measures to control crop pest & diseases can be taken to avoid their spread. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice (Mundakan) Panicle emergence stage	Sheath Blight	To control sheath blight disease in rice, spray 4 gram of (Trifloxystrobin + Tebuconazole) mixture in ten litres of water.
Banana	Calcium Boron Deficiency	To control leaf curling in banana, due to calcium deficiency, apply 250g lime per banana plant. Boron deficiency is seen more during dry weather. Spray one gram borax per one litre of water.
Pepper	Slow wilt	To control the slow wilt disease in pepper, apply neem cake at the base of pepper plants. Also apply 25 grams of the bio control agent <i>Pisilomyces lilacinus</i> at the base of each plant. If the attack is severe drench the basin of each plant with three gram Copper Oxy Chloride (COC) per one litre of water.

Animal Husbandry- Care should be taken to avoid ecto-parasitic infestation. Ensure sufficient supply of clean drinking water to the animals.