



Agromet Advisory Service bulletin 52/17, Friday dt. 29-12-2017

(Issued jointly by IAAS Vellanikkara)





ERNAKULAM

CURRENT SYNOPTIC SITUATION: A cyclonic circulation lies over Comorin area & neighbourhood and extends between 1.5 & 2.1 km above sea level.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 29.12.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

DISTRICT: ERNAKULAM

ENSEMBLE FCST

PARAMETERS	30.12.2017	31.12.2017	01.01.2018	02.01.2018	03.01.2018
Rainfall (mm)	0	0	0	0	0
Max Temperature (deg	32	32	33	33	33
Min Temperature (deg C)	23	23	24	24	24
Total cloud cover (octa)	4	3	3	3	2
Max Relative Humidity (%)	80	80	80	80	80
Min Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	6	6	6	6	6
Wind direction (deg)	230	230	230	230	230

Agro Meteorological Advisories:

gro Meteorological Advisories.				
Crop	Crop stage / Pest/Disease	Advisories		
Paddy (Jyothi, Kanchana)	Rice bug	Apply neem based pesticides (3ml per one liter of water). Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.		
Coconut	White fly	To control white fly, spray the mixture of 2% neem oil emulsion with 20 g of biocontrol agent (Verticillium) on lower sides of leaves. Avoid application of chemicals to save the biocontrol agents.		
Banana	Pseudostem weevil	I holes on pseudo stem will be drenched with chlorovritos 7 ml or		
Vegetables (Brinjal)	Shoot and Fruit borer	Shoot and fruit borer affected parts in brinjal can be removed. If severe infestation found, neem seed extract (5%) shall be sprayed. Sucking pest attacks like curling of leaf and leaf tips, yellowing can be prevented by spraying 2% neem oil garlic emulsion. Neem based other insecticides can also be applied intermittently.		
Animal Husbandry	If respiratory distress, fever or lameness are noticed in animals, consult a veterinary surgeon immediately. (CAADECS, KVASU, Mannuthy)			