



## Agromet Advisory Bulletin for the District, Kannur

(Valid from 02.10.2021 to 06.10.2021)

(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)



Bulletin Number: Pilicode/Knr-79/2021

Date: 01/10/2021

### A. Weather Summary of preceding five days





Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
59.8	29.2 – 32.5	24.1 – 25.5	80 – 96	00 – 08





### B. Weather forecast for next five days

Parameters	02-10-2021	03-10-2021	04-10-2021	05-10-2021	06-10-2021
Rainfall, mm	10	10	10	12	10
Max. Temp, °C	32	32	32	32	32
Min. Temp, °C	25	25	25	25	25
Max. Relative Humidity, %	96	96	96	96	96
Min. Relative Humidity, %	80	80	80	80	80
Wind speed, km/h	14	10	10	8	10
Wind direction, degrees	270	270	200	200	150
Total cloud cover, octa	8	8	8	8	8

### C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
<b>General conditions</b>			<b>Light to moderate rainfalls are very likely to occur on five days</b>
			<p>The sky will be cloudy or partially cloudy. Low night temperature and high day temperature will be experienced. Hence there will be large difference between day time temperature and night temperature. The atmospheric humidity also will be high.</p> <p><b>As per extended range forecast, cumulative normal rainfall is likely.</b></p> <p>Give popping to slender stemmed and easily lodging crops like banana, vegetables etc.</p> <p>The weather is favorable for the spread of fungal diseases like bud rot of coconut, mahali in arecanut, quick wilt in black pepper, rhizome rot diseases in ginger and turmeric.</p> <p>Take care while drying the harvested produces like rubber, cashew nut, copra etc., directly under the sun.</p>

Paddy second crop (Mundakan season)	Transplanting	Dip the roots of the seedlings in slurry of pseudomonas (20g pseudomonas/litre of water) for 30 minutes just before transplanting. This will help the seedlings to grow vigorously.	
Cool season vegetables	Land preparation	Lime or dolomite should be applied depending on soil acidity and incorporate thoroughly with the soil at least 14 days before the application of chemical fertilizers. This will not only reduce the acidity but also control the soil borne diseases.	
Coconut	All stages	<p style="text-align: center;">Leaf rot of coconut</p> 	<p>Remove and destroy the dried inflorescence and fallen nuts. Harvest all the matured nuts. After the harvest spray 1% Bordeaux Mixture or Copper oxy chloride (@3g/Litre) on the leaves and developing inflorescences and nuts</p> <p>After two weeks spray borax also on leaves @ 3g/l.</p>
Coconut	Bearing stage	<p style="text-align: center;">Nut rot</p> 	
Black pepper	All stages	<p style="text-align: center;">Gall midge</p> 	Spray dimethoate (@2ml/litre of water)
Ginger	All stages	<p style="text-align: center;">Rhizoctonia leaf blight</p> 	Spray carbendazim (@2g/litre of water)

Turmeric	All stages	<p>Shoot borer</p> 	Spray dimethoate (@2ml/litre of water)
Banana	Various stages of growth	<p>Panama wilt</p> 	<p>1. Prophylactic measures: Soil application of Pseudomonas @ 20g/litre of water. Dipping of suckers in carbendazim (@2g/litre of water) solution for 20 minutes before planting 2. If disease appears Crop sanitation: Eradication of infected plants and application of lime in the pits (@1kg/pit Drenching with carbendazim, @2g/L of water Spraying propiconazole @1ml/litre of water on leaves of the surrounding plants.</p>
Banana	Various stages of growth	<p>Sigatoka leaf spot</p> 	Cut and burn all severely affected leaves. Spray propiconazole (@2ml/Litre of water)r
Cashew	Flowering stage	<p>Tea mosquito bug</p> 	Spray Quinalphos @ 2ml/L.