




## Agromet Advisory Bulletin for the District, Kannur

(Valid from 06.10.2021 to 10.10.2021)





Bulletin Number: Pilicode/Knr-80/2021 Date: 05/10/2021

## A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
87.7	31.2 – 32.4	23.5 - 23.5	79.0 – 93.7	00 - 08

## B. Weather forecast for next five days

Parameters	06-10-2021	07-10-2021	08-10-2021	09-10-2021	10-10-2021
Rainfall, mm	7	8	10	10	5
Max. Temp, °C	32	32	32	32	32
Min. Temp, °C	25	25	25	25	25
Max. Relative Humidity, %	94	94	94	94	94
Min. Relative Humidity, %	81	81	81	81	81
Wind speed, km/h	5	3	6	5	6
Wind direction, degrees	230	180	180	200	200
Total cloud cover, octa	7	8	8	8	8

## C. Agrometeorological Advisories

Crop Stages		Problems	Agro-meteorological advisories	
General conditions		Light to moderate rainfalls are very likely to occur on five days.		
		Day-2: Heavy (7- 11 cm in 24 hours) rainfall is very likely to occur at isolated places in the district		
		The sky will be cloudy or partially cloudy. Low night temperature and high day temperature will be experienced. Hence there will be large difference between day time temperature and night temperature. The atmospheric humidity also will be high.		
		As per extended range forecast, cumulative normal rainfall is likely.		



		Complete the second dose crops quickly	of fertilizer applications to coconut like tree	
General conditions		If there are any crab holes /cracks on the bunds of the paddy fields that should be blocked by plastering with mud. Also remove all the weeds from the fields.		
		In plantation crops, water conservation measures should be adopted. Powdering the top soil uniformly spreading in the field, rain water harvesting pits (for which the bunds are strengthened by pineapple, vetiver etc.), cover cropping (especially with leguminous plants), coconut husk/coir pith drenching etc. will help to conserve the post monsoon rain water for a long period of time in the fields.		
Vegetables	Transplanting/ sowing	While preparing the land, incorporate lime @ 4kg/cent to the soil. Use Trichoderma enriched farm yard manure/compost. This will check the spread of wilt diseases.		
		Before transplanting, dip the roots of the seedlings in slurry of pseudomonas (20g pseudomonas/litre of water) for 30 minutes. This will help the seedlings to grow vigorously.		
		Also the seedlings can be sprayed with diluted pseudomonas culture solution (@20ml dissolved in one litre of water).		
Black pepper	All stages	Foot rot	As prophylactic measure, apply 150 gram of Trichoderma enriched neem cake - cow dung mixture in the basins of the vines and incorporate thoroughly with the soil.	
			If disease already appeared, drench soil in the plant basins with Redomil 0.2% (2g/litre of water). Spray the same on the leaves also.	
Ginger	All stages	Rhizoctonia leaf blight	Spray carbendazim (@2g/litre of water)	
Turmeric	All stages	Shoot borer	Spray dimethoate (@2ml/litre of water)	



Banana	Various stages of growth	Panama wilt	1. Prophylactic measures:  Soil application of Pseudomonas @ 20g/litre of water.  Dipping of suckers in carbendazim (@2g/litre of water) solution for 20 minutes before planting  2. If disease appears  Crop sanitation: Eradication of infected plants and application of lime in the pits (@1kg/pit Drenching with carbendazim, @2g/L of water Spraying propiconazole @1ml/litre of water on leaves of the surrounding plants.
Banana	Various stages of growth	Sigatoka leaf spot	Cut and burn all severely affected leaves. Spray propiconazole (@2mlLitre of wate)r
Cashew	Newly planted seedlings	Tea mosquito attack and associated fungal diseases	Prophylactic spray of combination Bordeaux mixture (1% strength) + quinalphos (2ml/litre of the Bordeaux mixture )

Sd/-Nodal Officer, GKMS Project, RARS Pilicode