



# Agromet Advisory Bulletin for the District, Kannur

(Valid from 23.12.2020 to 27.12.2020)

(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)



<b>Bulletin Number: Pilicode/Knr-102/2020</b>	<b>Date: 22/12/2020</b>
---	-------------------------


## A. Weather Summary of preceding five days







Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
0.0	34.2 – 34.6	22.4-25.2	57–78	04 -12

## B. Weather forecast for next five days

Parameters	23.12.2020	24.12.2020	25.12.2020	26.12.2020	27.12.2020
Rainfall, mm	0	4	2	2	0
Max. Temp, °C	34	34	34	34	34
Min. Temp, °C	23	23	23	23	23
Max. Relative Humidity, %	78	78	78	78	78
Min. Relative Humidity, %	57	57	57	57	57
Wind speed, km/h	16	14	14	18	16
Wind direction, degrees	200	250	270	230	250
Total cloud cover, octa	1	4	4	6	1

## C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
<b>General conditions</b>			<p><b>Isolated rainfalls and strong wind may occur.</b></p> <p><b>Prop bananas and vegetables</b></p> <p><b>Copra and Rubber, Arecanut etc should not be left in the drying yards during night.</b></p>
Bitter gourd	All stages	<p>Fruit fly</p> 	<p>Setting up of pheromone trap(Cue lure @1/15 cent)</p> <p>Spot application of 10 % jaggery containing 0.1 % malathion@1 spot/40 m<sup>2</sup> on underside of leaves at fortnight intervals.</p>

Chilli	Active growing stage	<b>Thrips and Mites</b> 	Spray 'spiromesifen' (@1ml in litre of water)
Cabbage	All stages	<b>Black rot</b> 	Spray Streptocyclin 300ppm. Spray Pseudomonas 2% solution weekly once after head initiation.
Chilli	All stages	<b>White fly</b> 	Spray Thiamethoxam @ 2gm/10litre of water.
Cowpea	Growing stage	<b>Wilt</b> 	Spray Contaf (@ 2ml/litre)
Cucurbits	All stages	<b>Downy mildew</b> 	As a prophylactic measure apply 'Mancozeb'(@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) on both surfaces of the leaves, thrice at 15 days interval.or spraying of pyraclostrobin @.5gm/L.
Pumpkin	All stages	<b>Pumpkin beetle</b> 	Apply malathion 50 EC @ 3 ml / litre of water or malathion 50 EC @ 2 ml + neem oil 4ml per litre of water.  After irrigation, drench the soil with malathion 50 EC @ 3 ml / litre of water. After the application of insecticide, with hold irrigation on next two days.