

## **Agromet Advisory Bulletin for the District, Kannur**

(30.11.2022 to 04.11.2022)



(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)

Bulletin Number: Pilicode/Knr-95/2022 Date: 29/11/2022

## A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
0.2	32.3 - 33.7	24.6 – 25.8	72.9 – 90.4	00 – 6.0

## B. Weather forecast for next five days

Parameters	30-11-2022	01-12-2022	02-12-2022	03-12-2022	04-12-2022
Rainfall, mm	2	0.1	0	0	0.1
Max. Temp, °C	33	33	33	33	33
Min. Temp, °C	23	23	23	23	23
Max. Relative Humidity, %	90	90	90	90	90
Min. Relative Humidity, %	73	73	73	73	73
Wind speed, km/h	2	2	2	2	2
Wind direction, degrees	270	270	250	270	250
Total cloud cover, octa	4	2	0	2	8

## C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories		
	Isolated light rainfalls very likely to occur **				
General conditions	Isolated light rainfalls very likely to occur.  Low night temperature (minimum 24°C) and high day temperature (maximum 33°C) will be experienced. Hence there will be large difference between day time temperature and night temperature.  The atmospheric humidity also will be high (between 76 and 90%).				
General Recommen dations	Rain: Keep vigilance while sun drying perishable goods, rice, copra and rubber like products. Provide propping to tender stem crops like banana, tomato etc. Give popping to slender stemmed and easily lodging crops like banana, vegetables etc.				
	Protect newly planted seedlings of the crops from the bright sunshine from southern side, by smearing with banana/coconut leaves.				
	Maintain hygiene conditions in crop fields. Infected and fallen nuts, leaves and tree parts should be removed from the fields and burnt.				
	Fungal and bacterial diseases like blight, leaf spots and wilt may spread in crops. Keep vigilance. Take control measures in the beginning stage of diseases.				

Paddy	All stages	Bacterial leaf blight	Spray supernatant solution of the cow dung water mixture (Dispense 20g cow dung in 1 L of water and filter the solution through muslin cloth). Place bags containing bleaching powder (2kg/acre) in the paddy field at the entry face of running water into the field  If the infection is severe spray Streptocyclin 30g/200 L of water for 1acre.
Coconut	Bearing stage	Mahali	Harvest matured nuts. After that spray 1% Bordeaux mixture to the affected nuts/bunches
Cucurbit Vegetables	Planting	Downy mildew	As a prophylactic measure apply 'Mancozeb'(@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) on both surfaces of the leaves, thrice at 15 days interval. Remove and burn out the infested leaves.
Cucurbitaceo us Vegetables	Flowering and fruit setting	Melon fruit flies	Keep pheromone traps.  Spray Malathion:- Mix 2 ml of Malathion in one litre of water. Dissolve 10g of jaggery in the solution to attract the flies. Spray this solution on lower sides of the leaves
Ash gourd/Cucu rbits	Fruit bearing	Fruit rot	Spray Tebuconazole (@1ml/litre)
Brinjal	Flowering and fruit setting	Shoot and Fruit borer	Remove and destroy affected fruits and shoots. Spray chlorantraniliprole (coragen 3ml per 101 of water) 2-3 times at weekly interval.

Mango	Fruit setting stage	Mango fruit flies	Collect and destroy the fallen fruits by taking deep pits atleast 60 cm depth.  Set up pheromone trap (methyl eugenol trap) @ 1 trap/15 cents.
Nutmeg	Fruiting	Leaf/Fruit spots	Spray copper oxychloride (@2g/litre)

\*\* Warning colour codes of rainfall (for disaster management)

Warning (Take actions)   Alert (Be prepared)   Watch (Be updated)   No warning	
	g (No actions)

Sd/-Nodal Officer, GKMS Project, RARS Pilicode