

आजादी
का
अमृत महोत्सव



Agromet Advisory Bulletin
Kasaragod District
(Valid from 20.04.2022 to 24.04.2022)



Agromet Advisory Bulletin for the District, Kasaragod

(Valid from 20.04.2022 to 24.04.2022)

(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)



Bulletin Number: Pilicode/Ksd-31/2022

Date:19/04/2022

A. Weather Summary of preceding five days

| Rainfall, mm | Max. temp., °C | Min. temp., °C | R. H., % | Wind speed, Km/h |
|--------------|----------------|----------------|----------|------------------|
| 82.5 | 32-35 | 24.5-26.5 | 77-94 | 0.0-4.0 |




B. Weather forecast for next five days



| Parameters | 20-04-2022 | 21-04-2022 | 22-04-2022 | 23-04-2022 | 24-04-2022 |
|---------------------------|------------|------------|------------|------------|------------|
| Rainfall, mm | 2 | 1 | 0 | 0 | 0 |
| Max. Temp, °C | 35 | 35 | 35 | 35 | 35 |
| Min. Temp, °C | 26 | 26 | 26 | 26 | 26 |
| Max. Relative Humidity, % | 77 | 77 | 77 | 77 | 77 |
| Min. Relative Humidity, % | 60 | 60 | 60 | 60 | 60 |
| Wind speed, km/h | 6 | 6 | 6 | 6 | 6 |
| Wind direction, degrees | 250 | 250 | 250 | 250 | 250 |
| Total cloud cover, octa | 4 | 5 | 4 | 4 | 4 |

C. Agrometeorological Advisories

| Crop | Stages | Problems | Agro-meteorological advisories |
|---------------------------|---|----------|--------------------------------|
| General conditions | Isolated light to moderate rainfalls with thunderstorm and heavy wind | | |
| | <p>The sky will be cloudy or partially cloudy. High atmospheric temperature and relative humidity will be experienced.</p> <p>There will be light to moderate rainfalls with lightning and thunderstorm in isolated places.</p> | | |

| | | |
|--|--|--|
| <p><u>General Recommendations</u></p> | <p>Harvest all the ripened fields quickly. Do not heap the harvested products in the field itself. Take care while drying the harvested rice directly under the sun, especially during the afternoon periods.</p> <p>Take care while drying the harvested products like rice, rubber, copra, pulses, vegetable seeds, etc directly under sun.</p> <p>Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc. Clean the drainage channels in crop lowlands to enable proper drainage of excess water in case of exigencies if any.</p> <p>Keep animal feeds, seeds, fertilizers etc. in termite free and moist free rooms on elevated platforms coated with wooden pannels.</p> <p>Summer rains have prime role in coping up drought. Hence maximum water harvesting should be ensured in the fields. Clean the rain pits. Cover the soil with dried leaves, especially the basins of crops. The opened tree basins which are partially filled after fertilizer application, can act as very good water harvesting structures. Divert the runoff water to such tree basins by drawing furrows.</p> <p>In rice fallows, organic manure seeds (Daincha, sunhemp, cowpea etc) can be sown during this time. These can be harvested at its 45 days of growth, just before flowering. This will not only help in fixing nitrogen into the soil, but also control the weeds growth in the fields. If there is sufficient soil moisture at the time of harvest, incorporate them into the soil through deep ploughing. Other wise use it for preparation of composts.</p> <p>Avoid activities under open field conditions during 12 AM to 3 PM on the days with high temperature. Also avoid activities under open conditions while there is lightning and heavy winds. Also do not expose livestock to such unfavorable weather conditions.</p> | |
| <p>Paddy (Late sown Mundakan crop)</p> | <p>Harvesting</p> | <p>Harvest all the ripened fields quickly. Do not heap the harvested products in the field itself. Take care while drying the harvested rice directly under the sun, especially during the afternoon periods.</p> |
| <p>Paddy (Viruppu: First crop season)</p> | <p>Land preparation for broadcasting</p> | <p>Summer showers are predicted. Immediately after attaining sufficient moisture level in soil, land preparations can be started in places where broadcasting is preferred.</p> <p>In rice fallows where transplanting is practiced during the first crop, organic manure seeds (Daincha, sunhemp, cowpea etc) can be sown during this time. These can be harvested at its 45 days of growth, just before flowering. This will not only help in fixing nitrogen into the soil, but also control the weeds growth in the fields. If there is sufficient soil moisture at the time of harvest, incorporate them into the soil through deep ploughing. Other wise use it for preparation of composts.</p> |
| <p>Black Pepper</p> | <p>Bearing</p> | <p>High intensity rainfalls may cause non uniform pollination as the pollen grains might have taken away quickly by the high velocity of the falling raindrops. To ensure uniform pollination in the spikes, lightly spray water on rain free days.</p> |

| | | | |
|--------------|---------------|--|---|
| Coconut | All stages | <p>Rhinoceros beetle</p>  <p>Dwarf varieties and young palms are more vulnerable to the attack.</p> | <p>Take out the beetles from the attacked palm crown using beetle hook. Swab the hole made by the beetles with Bordeaux paste to prevent entry of fungus through the cut surface.</p> <p>Old fishnets can be reused for controlling the pest. Cut the nets into pieces of size 1m x 0.5m. Keep the middle portion of the net, lengthwise into the axils of 3rd and 4th leaf whirls of the palm. Push the net firmly with small stones into the axils. The hanging sides of the net should be kept open. The beetles will be trapped in the nets and get destroyed. Care should be taken to move the nets to upper leaf axils accordingly while fresh leaves are emerging.</p> |
| Coconut | All stages | <p>Bud rot</p>  | <p>Detection of disease at its early stage will help to adopt efficient corrective measures. Cut and remove the affected tissues from the crown and apply Bordeaux paste. After that cover the cut surface with polythene sheets to protect it from rain falls until new leaf emerges.</p> <p>Burn the removed tissues immediately.</p> <p>As a prophylactic measure apply Trichoderma cakes of spray 1% Bordeaux mixture into the axils of top leaves of the surrounding palms</p> |
| Black pepper | All stages | <p>Foot rot</p>  | <p>As prophylactic measure, apply 150 gram of Trichoderma enriched neem cake - cow dung mixture in the basins of the vines and incorporate thoroughly with the soil.</p> <p>If disease already appeared, drench soil in the plant basins with Redomil 0.2% (2g/litre of water). Spray the same on the leaves also.</p> |
| Areca nut | Bearing Stage | Mahali disease | Apply Bordeaux mixture (1%) to the bunch and leaves. |

| | | | |
|--------|--------------------------|---|---|
| Banana | Various stages of growth | <p style="text-align: center;">Sigatoka leaf spot</p>  | <p>Cut and burn all affected leaves.</p> <p>Spray Tilt or Contaf (@ 2ml per litre)</p> |
| Banana | All stages | <p style="text-align: center;">Pseudo stem weevil</p>  | <p>Adopt good field sanitation.</p> <p>Remove the affected parts and burn it using kerosene.</p> <p>Apply Beauveria bassiana @ 20 g l⁻¹ at 5, 6 and 7 month after planting, into the leaf axils</p> <p>If attacked holes and yellowing are seen, apply Fipronil 0.015% (3 ml l⁻¹) into the leaf axils</p> |

Sd/-
Nodal Officer,
GKMS Project, RARS Pilicode