

Bulletin Number: Pilicode/Ksd-68/2020

Date: 25/08/2020



A. Weather Summary of preceding five days





Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
36.6	28.0 – 31.0	22.5 – 24.0	75– 96	01 -02

B. Weather forecast for next five days

Parameters	26.08.2020	27.08.2020	28.08.2020	29.08.2020	30.08.2020
Rainfall, mm	0	0	0	0	0
Max. Temp, °C	31	31	31	31	31
Min. Temp, °C	24	24	24	24	24
Max. Relative Humidity , %	95	95	95	95	95
Min. Relative Humidity, %	68	68	68	68	68
Wind speed, km/h	16	16	16	8	10
Wind direction, degrees	320	320	320	290	290
Total cloud cover, octa	5	4	5	6	6

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
General conditions		Rainfalls are unlikely on coming days.	
Paddy	Grain filling stage	<p style="text-align: center;">Brown spot</p> 	Foliar spraying of Hexaconazole (@ 2 ml/liter water)
Coconut	All stages	<p style="text-align: center;">Bud rot</p> 	<p>As a preventive measure, small perforated sachets containing 2 g of mancozeb may be tied to the top leaf axil. When rains, a small quantity of the fungicide is released from the sachets to the leaf base and thus protecting the palm.</p> <p>Spray 1% Bordeaux mixture on the crown of palms.</p>

Areca nut	All stages	<p>Mahali disease</p> 	<p>Apply Bordeaux mixture (1%) to the bunch and leaves.</p>
Black pepper	All stages	<p>Quick wilt</p> 	<p>Spray Redomil (@2g/litre of water) or copper oxy chloride (@3g/litre of water)</p> <p>Drench the soil in the basins with the same preparation (@ 5 litre per vine) as mentioned above.</p>
Cowpea	Pod development stage	<p>Pod bug</p> 	<p>Spray Takumi @ 2 gm/ 10 L.</p>
Okra	All stages	<p>Shoot and Fruit borer</p> 	<p>Spray neem oil emulsion @ 5 %, at intervals of 15 to 20 days. Spraying with quinalphos 25 EC (2 ml per litre of water).</p>

Sd/-
Nodal Officer,
GKMS Project, RARS Pilicode