



Agromet Advisory Bulletin for the District, Kozhikode (Valid from 22.12.2021 to 26.12.2021)



(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)

Bulletin Number: Pilicode/Kkd-102/2021 Date: 21/12/2021

A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
00	33.6 - 34.6	23.0 - 25.2	59.6 - 87.2	00 - 06

B.Weather forecast for next Three days

Parameters	22-12-2021	23-12-2021	24-12-2021	25-12-2021	26-12-2021
Rainfall, mm	110	-0		0	0
Max. Temp, °C		34	34	34	34
Min. Temp, °C	23	23	23.00	23	23
Max. Relative Humidity, %	85	85	85	85	85
Min. Relative Humidity, %	60	60	60	60	60
Wind speed, km/h	5 🕈		4	5	6
Wind direction, degrees	270	270	270	270	270
Total cloud cover, octa	2			1	1

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
<u>General</u> <u>conditions</u>	 will be experiencender in the imperature Fungal diseases la control measures In plantation cropuniformly spreaded strengthened by presented 	ed. Hence there will be large b. The atmospheric humidity also ike blight, leaf spots and wil lt in the beginning stage of diseas os, water conservation measure ling in the field, rain water pineapple, vetiver etc.), cover of r pith drenching etc. will help t	may spread in crops. Keep vigilance. Take



Paddy	Grain filling	Rice bug	Strict vigilance is necessary at milky stage. Keep the field and bunds free of weeds and grasses. If attack is seen apply fish amino acid @ 20 ml/litre.
Coconut Black pepper	All stages All stages	Rhinoceros beetle Warf varieties and young palms are more vulnerable to the attack. Pollu disease	Take out the beetles from the attacked palm crown using beetle hook. Swab the hole made by the beetles with Bordeaux paste to prevent entry of fungus through the cut surface. Old fishnets can be used for controlling the pest. Cut the nets into pieces of size 1 m x 0.5m. Keep the middle portion of the net, lengthwise into the axils of 3 rd and 4 th leaf whirls of the palm. Push the net firmly with small stones into the axils. The hanging sides of the net should be kept open. The beetles will be trapped in the nets and get destroyed. Care should be taken to move the nets to upper leaf axils accordingly while fresh leaves are emerging. Spray hexaconazole (@ 2ml/litre of water)
Fobber			The second
Cashew	Seedlings (up to four years)	Tea mosquito attack and associated fungal diseases	Prophylactic spray of combination of Bordeaux mixture (1% strength) + quinalphos (2ml/litre of the Bordeaux mixture) If die back is seen apply combination of hexaconazole and malathione (@ hexaconosozole 1ml + malathione1ml per litre of water)



Jack	Various stages	Fruit fall due to fungal infections	Spray Bordeaux mixture (1% strength)
Mango	Flowering stage	Mango hoppers	Spray Azadirachtin (@5ml/litre Azadirachtin 3000pmm)
Mango	Different stages	Anthracnose disease	Spray copper oxy chloride or hexaconazole (@ 2ml/litre of water)
Bitter gourd	All stages	Downy mildew	As a prophylactic measure apply 'Mancozeb'(@ 2g/l of water). If disease appeared, spray Saaf (2g/litre of water)
Bovines		e lumpy skin and foot and uth diseases	Adopt vaccination. Keep cattle sheds and its surroundings hygienic.
Goats	Р	PR disease	Adopt vaccination. Keep cattle sheds and its surroundings hygienic