



# Agromet Advisory Bulletin for the District, Kozhikode

(Valid from 05.12.2020 to 09.12.2020)

(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)



Bulletin Number: Pilicode/Kkd-97/2020

Date: 04/12/2020


## A. Weather Summary of preceding four days






Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
1.2	30.6 – 34.4	23.8 – 24.9	63– 96	00-12

## B. Weather forecast for next five days

Parameters	05.12.2020	06.12.2020	07.12.2020	08.12.2020	09.12.2020
Rainfall, mm	10	10	10	10	12
Max. Temp, °C	30	30	30	30	30
Min. Temp, °C	24	24	24	24	24
Max. Relative Humidity, %	88	88	88	88	88
Min. Relative Humidity, %	63	63	63	63	63
Wind speed, km/h	4	6	2	10	6
Wind direction, degrees	290	250	340	290	250
Total cloud cover, octa	8	8	8	8	8

## C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
<b>General conditions</b>		<p><b>Rainfalls and strong wind may occur.</b></p> <p><b>Prop bananas and vegetables</b></p> <p><b>Copra and Rubber, Arecanut etc should not be left in the drying yards during night.</b></p>	
Coconut	All stages	<p>Basal rot</p> 	<p>Drench the basin with Hexaconazole (25 ml in 25 L of water).</p> <p>After one week apply lime @ 2 kg per palm.</p> <p>Apply Potash @ 1 kg per palm three weeks after lime application to improve the health of palm, in addition to the regular dose.</p> <p>Avoid flood irrigation to prevent the spread of pathogen.</p>

Cabbage	All stages	<p>Black rot</p> 	<p>Spray Streptocyclin 300ppm. Spray Pseudomonas 2% solution weekly once after head initiation.</p>
Chilly	All stages	<p>Bacterial wilt</p> 	<p>Keep the field clean and spray antibiotic strepto cycline 1 capsule per 2 litre or drench the field with COC 2gm/L.</p>
Cowpea	Growing stage	<p>Wilt</p> 	<p>Spray Contaf (@ 2ml/litre)</p>
Cucurbits	All stages	<p>Downy mildew</p> 	<p>As a prophylactic measure apply 'Mancozeb' (@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) on both surfaces of the leaves, thrice at 15 days interval.or spraying of pyraclostrobin @.5gm/L.</p>
Pumpkin	All stages	<p>Pumpkin beetle</p> 	<p>Apply malathion 50 EC @ 3 ml / litre of water or malathion 50 EC @ 2 ml + neem oil 4ml per litre of water. After irrigation, drench the soil with malathion 50 EC @ 3 ml / litre of water. After the application of insecticide, with hold irrigation on next two days.</p>

Sd/-  
Nodal Officer,  
GKMS Project, RARS Pilicode