

Agromet Advisory Bulletin for the District, Kozhikode (Valid from 18.10.2023to 22.10.2023)



(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode& India Meteorological Department)

Bulletin Number:Pilicode/Kkd-83/2023 Date:17/10/2023

A.Weather Summary of preceding five days

| Rainfall, mm | Max. temp., °C | Min. temp., °C | R. H., % | Wind speed, Km/h |
|--------------|----------------|----------------|----------|------------------|
| 25.7 | 31.8 - 34.0 | 23.6 - 26.2 | 72 - 91 | 00 - 08 |

B.Weather forecast for next five days

| Parameters | 18-10-2023 | 19-10-2023 | 20-10-2023 | 21-10-2023 | 22-10-2023 |
|---------------------------|------------|------------|------------|------------|------------|
| Rainfall, mm | 4 | 5 | 4 | 4 | 2 |
| Max. Temp, °C | 33 | 34 | 34 | 34 | 34 |
| Min. Temp,°C | 25 | 25 | 25 | 25 | 25 |
| Max. Relative Humidity, % | 95 | 95 | 95 | 92 | 92 |
| Min. Relative Humidity, % | 70 | 60 | 60 | 60 | 65 |
| Wind speed,km/h | 5 | 6 | 5 | 5 | 5 |
| Wind direction, degrees | 270 | 290 | 270 | 250 | 250 |
| Total cloud cover, octa | 8 | 8 | 8 | 7 | 8 |

C. Agrometeorological Advisories

| Crop | Stages | Problems | Agro-meteorological advisories | | |
|-----------------------------------|---|----------|--------------------------------|--|--|
| | Isolated heavy rainfalls ** | | | | |
| <u>General</u> conditions | The sky will be cloudy or partially cloudy. Low night temperature and high temperature will be experienced. Hence there will be distinct difference between time temperature and night temperature. The atmospheric humidity also will be high. | | | | |
| conditions | There will be isolated heavy rainfalls (up to 115.5 mm within a time span of 24 hour most places are very likely on October 17 th . | | | | |
| | There will be light rainfalls (from 2.5mm to 15.5 mm within a time span of 24 hours) in most places from October 18 th to October 21 th . | | | | |
| <u>General</u> | Keep all drainage channels opened in the cultivated areas | | | | |
| <u>Recommen</u> <u>dations</u> | Transport perishable produces like rubber, copra, pulses, vegetables, seeds, etc. and animal feeds and fertilizers from open places to store houses. Collect and store sufficient feeds for the livestock. Keep the seeds fertilizers and cattle feeds on moisture free and elevated plant forms. | | | | |
| | Neither keep the livestock in open places, nor allow them for grazing. They should be | | | | |

| | sheltered under strong roofs. | | | | |
|---------------------------|------------------------------------|---|---|--|--|
| | Postpone activities lik favorable. | e fertilizer application, sowing etc. till climatic conditions become | | | |
| Cool season vegetables | Land preparation | Lime or dolomite should be applied depending on soil acidity and incorporate thoroughly with the soil at least 14 days before the application of chemical fertilizers. This will not only reduce the acidity but also control the soil borne diseases. | | | |
| Coconut | All stages | Bud rot | Detection of disease at its early stage will help to adopt efficient corrective measures. Cut and remove the affected tissues from the crown and apply Bordeaux paste. After that cover the cut surface with polythene sheets to protect it from rain falls until new leaf emerges. Burn the removed tissues immediately. As a prophylactic measure spray 1% Bordeaux mixture into the axils of top leaves of the surrounding palms | | |
| Coconut | All stages | Rhinoceros beetle Warf varieties and young palms are more vulnerable to the attack. | Take out the beetles from the attacked palm crown using beetle hook. Swab the hole made by the beetles with Bordeaux paste to prevent entry of fungus through the cut surface. Old fishnets can be reused for controlling the pest. Cut the nets into pieces of size 1m x 0.5m. Keep the middle portion of the net, lengthwise into the axils of 3 rd and 4 th leaf whirls of the palm. Push the net firmly with small stones into the axils. The hanging sides of the net should be kept open. The beetles will be trapped in the nets and get destroyed. Care should be taken to move the nets to upper leaf axils accordingly while fresh leaves are emerging. | | |
| Coconut | All stages | Yellowing of coconut | Incorporate lime 1kg + dolomite 1kg (per palm) to the basin of the palms. After two weeks, apply borax @ 200g/palm and zinc sulphate @ 100g/palm and magnesium sulphate @500g/palm and potash @ 2kg/palm | | |

| growthin the basins of the crop. As a prophylactic measure spray Psedudmonas (@ 20g/lire) at fortnight intervals.CashewPre-bearing stagesTea mosquito bugCashewPre-bearing stagesTea mosquito bugPoultryAny stageNewcastle Disease Image of the composition of the comp | Black pepper | All stages | Foot rot | As prophylactic measure, apply 150 gram of Trichoderma enriched neem cake - cow dung mixture in the basins of the vines and incorporate thoroughly with the soil.If disease already appeared, drench soil in the plant basins with Redomil 0.2% (2g/litre of water). Spray the same on the leaves also. |
|---|--------------|--------------------|--------------------|--|
| Calle II Field and generation Poultry Any stage Poultry Any stage Newcastle Disease Image: Image Ima | Banana | - | Sigatoka leaf spot | As a prophylactic measure spray Pseudomonas (@ 20g/litre) at fortnight intervals. If disease appears:-Cut and burn all affected leaves. Spray Carbendazim and Mancozeb (@ 2ml per litre) alternatively |
| GoatAll stagesAllergic diarrheaTo prevent dehydration and electrolyte losses give Ringer's lactate solution intravenously in consultation with the | Cashew | Pre-bearing stages | Tea mosquito bug | The fungal biological control agent, Beavaria bassiana can be sprayed @ 20g/L Or 2) Spray 1% Bordeaux mixture mixed with quinalphos (2ml/litre of Bordeaux |
| GoatAll stagesAllergic diarrheaTo prevent dehydration and electrolyte losses give Ringer's lactate solution intravenously in consultation with the | Poultry | Any stage | Newcastle Disease | coughing, drooping wings, dragging legs, twisting of the head and neck, circling, in appetence, complete paralysis. |
| losses give Ringer's lactate solution intravenously in consultation with the | | | | condition for birds and chicken coops. Provide vaccinations with Live B1 and La Sota strains administrated in drinking water intranasally or |
| ** Warning colour codes of rainfall (for disaster management) | | | | To prevent dehydration and electrolyte losses give Ringer's lactate solution intravenously in consultation with the nearest veterinary doctor. |

| Warning (Take actions) | Alert (Be prepared) | Watch (Be updated) | No warning (No actions) |
|------------------------|---------------------|--------------------|-------------------------|

Sd/-Nodal Officer, GKMS Project, RARS Pilicode