



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



**PALAKKAD DISTRICT**

AAS 04/2019(7), Tuesday

22.01.2019

**CURRENT SYNOPTIC SITUATION:** A Low Pressure Area lies over Equatorial Indian Ocean & adjoining south East Bay of Bengal with associated cyclonic circulation extending upto 3.1 km above mean sea level. A cyclonic circulation extending upto 1.5 km above mean sea level lies over south Madhya Pradesh and neighbourhood. A trough runs from the above cyclonic circulation to South Interior Karnataka across Madhya Pradesh and North interior Karnataka at 1.5 km above mean sea level.

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	26.01.2019	27.01.2019	28.01.2019	29.01.2019	30.01.2019
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	35	35	35	35	35
Minimum temperature (°C)	22	22	22	22	22
Total cloud cover (octa)	4	5	6	6	4
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	35	35	35	35	35
Wind speed (kmph)	8	8	8	12	12
Wind direction (deg)	250	250	250	70	70

**No rainfall is forecasted for coming five days in the district by India Meteorological Department.**

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Crop stage	Pest/ Disease	Advisories
Paddy	Milky stage	Rice bug	There is a chance of appearance of rice bug in paddy. To control them, spray 2 ml malathion per one litre of water. Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.
Coconut	-	Red palm weevil	Coconut palms are seen affected by red palm weevil. Remove the affected parts from crown and apply one ml imidacloprid /one litre of water at spindle leaf area.
Vegetables (cowpea)	-	Aphids	There is a chance of aphids in cowpea. Apply 2% neemoil emulsion or apply 20 g verticillium lecanii per one litre of water in 10 days intervals. If aphid attack is severe, apply 3 ml Imidacloprid or 2 g Thiamethoxam per 10 litre of water.
Animal Husbandry	Mix 30g sodium bicarbonate and 1 teaspoon yeast in the feed daily to avoid digestive disorders associated with excessive acidity. (CAADECCS, KVASU, Mannuthy)		