



## Agromet Advisory Bulletin for PALAKKAD District

(Valid from 29.01.2025 to 02.02.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara  
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 04/2025, (8) Tuesday

Dated 28.01.2025

### A. Medium range weather forecast of IMD for next five days

Weather parameters	29.01.2025	30.01.2025	31.01.2025	01.02.2025	02.02.2025
Rainfall (mm)	0	0	0.1	0.1	0.1
Maximum temperature (°C)	33	33	33	33	32
Minimum temperature (°C)	23	23	23	23	24
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	40	40	40	40	40
Wind speed (kmph)	10	12	8	8	8
Wind direction (deg)	80	80	90	90	80
Total cloud cover (octa)	3	4	7	8	8

**Weather summary / Alert:** Since dry weather condition is prevails, irrigation of crops can be continued as per requirement. Normal to above normal rainfall is expected in Kerala.

### B. Agrometeorological Advisories for crop protection & management

#### General and Agrometeorological Advisories for Thrissur District

**General Advisory:** - Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Rice (Mundakan) Maturity stage	Lakshmi disease	There is a chance of Lakshmi disease in rice. To control this, spray 2 gram Copper hydroxide in one liter of water.
	Sheath Blight	To control sheath blight disease in rice, spray 4 gram of Trifloxystrobin + Tebuconazole mixture in ten litres of water.
Banana	Management	Since the wind speed may increase, give strong propping to banana plants.
Pepper	Slow wilt	To control the slow wilt disease in pepper, apply neem cake at the base of pepper plants. Also apply 25 grams of the bio control agent <i>Pisilomyces lilacinus</i> at the base of each plant. If the attack is severe drench the basin of each plant with three gram Copper Oxy Chloride (COC) per one litre of water.
Chilli	Thrips	There is a chance of thrips attack in chilli. Apply 2% neem oil+ garlic emulsion under the leaf surface at 10 days intervals. If the attack is severe, apply spiromecifen 8 ml per ten litre of water.

**Animal Husbandry:** Ensure sufficient supply of clean drinking water to the animals. Take measures for preventing mosquito breeding in the premises. (CAAECCS, KVASU, Mannuthy).