



Agromet Advisory Bulletin for PALAKKAD District

(Valid from 05.02.2025 to 09.02.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 05/2025, (10) Tuesday

Dated 04.02.2025

A. Medium range weather forecast of IMD for next five days

Weather parameters	05.02.2025	06.02.2025	07.02.2025	08.02.2025	09.02.2025
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	34	34	34	34	34
Minimum temperature (°C)	21	21	21	21	21
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	30	30	30	30	30
Wind speed (kmph)	4	4	4	4	4
Wind direction (deg)	70	70	70	70	70
Total cloud cover (octa)	4	3	3	3	3

Weather summary / Alert: Since dry weather condition is prevails, irrigation of crops can be continued as per requirement. Normal to above normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

General and Agrometeorological Advisories for Thrissur District

General Advisory: - Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Rice (Mundakan) Maturity stage	Bacterial Leaf Blight	To control and prevent Bacterial leaf blight in paddy, dissolve 20 g of cow dung in one litre of water and spray the supernatant liquid by mixing with 20g of pseudomonas. If the attack is severe spray 6 g of Streptocycline in 20 litre of water.
Bhindi	Jassid's	Due to increase in atmospheric temperature, the attack of Jassid's in Bhindi was increased. They can be controlled by applying 2% neem oil emulsion once in a week or by spraying 2 g thiamethoxam per ten litre of water.
Chilli	Thrips	There is a chance of thrips attack in chilli. Apply 2% neem oil + garlic emulsion under the leaf surface at 10 days intervals. If the attack is severe, apply spiromecifen 8 ml per ten litre of water.

Animal Husbandry: Ensure sufficient supply of clean drinking water to the animals. Take measures for preventing mosquito breeding in the premises. (CAAECCS, KVASU, Mannuthy).