



Agromet Advisory Service bulletin 14/16, Tuesday dt.29-03-2016

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

PALAKKAD

CURRENT SYNOPTIC SITUATION: A trough of low at mean sea level lies over Southwest Bay of Bengal off Sri Lanka coast.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 29.03.2016 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 30/03	DAY-2 31/03	DAY-3 01/04	DAY-4 02/04	DAY-5 03/04
DISTRICT : PALAKKAD					
Rainfall (mm)	0	0	0	0	0
Max Temperature (deg C)	40	40	40	40	40
Min Temperature (deg C)	27	27	27	27	27
Total cloud cover (octa)	5	6	4	4	4
Max Relative Humidity (%)	80	80	80	80	80
Min Relative Humidity (%)	40	40	40	40	40
Wind speed (kmph)	006	006	006	006	006
Wind direction (deg)	250	250	250	250	250

Agro Meteorological Advisories

Crop	Pest/Disease	Advisories
Donono	Cicatalra	There is a chance of sigatoka disease in banana. Spray Pseudomonas
Banana	Sigatoka	10- 20g per 1 liter of water. In case of severe attack apply SAAF @ 2 g in 1 liter of water
		2 g III 1 liter of water
		Burying of fresh or dried coconut husk around the palm is a
		desirable practice particularly for moisture retention. Mulch the
Coconut	Management	coconut basins with green/dry leaves. This help to add organic
		matter to the soil and to reduce the soil temperature. Drip irrigation
		may be practicised in coconut plantations.
		Due to the increase in atmospheric temperature, the attack of insects
		such as aphids, white flies, hoppers are more. As a result of this,
Vegetables	Mosaic	there is a chance of mosaic disease in vegetables, which can be
		controlled by applying 5 gms of Verticillium or Metarhyzium in 1
		litre of water. Add neem cake as manure. Neem based insecticides
		can be applied.