



Agromet Advisory Bulletin for PALAKKAD District

(Valid from 16.04.2025 to 20.04.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 15/2025, (30) Tuesday

Dated 15.04.2025

A. Medium range weather forecast of IMD for next five days

Weather parameters	16.04.2025	17.04.2025	18.04.2025	19.04.2025	20.04.2025
Rainfall (mm)	0.2	1	0.5	0.1	0.2
Maximum temperature (°C)	37	37	37	37	37
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	85	85	85	85	85
Minimum Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	2	2	2	8	8
Wind direction (deg)	200	250	230	250	230
Total cloud cover (octa)	8	8	8	8	8

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of light rainfall in the district. Thunderstorm accompanied by lightning and gusty winds is very likely to occur at one or two places over Kerala on 15th to 19th April 2025. Above normal rainfall is expected in Kerala.

Maximum temperatures are very likely to be around 37°C in 15th & 16th April 2025. Hot & Humid weather is very likely to prevail over the district except in hilly areas due to high temperature and humidity.

B. Agrometeorological Advisories for crop protection & management

General and Agrometeorological Advisories for Palakkad District

General Advisory: - Proper mulching can be followed to conserve soil moisture. Since air temperature is increasing, it is a desirable practice to bury fresh or dried coconut husk in the coconut basin to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Puncha rice (Dough stage)	Rice bug	There is a chance of appearance of rice bug in paddy. To control spray fish jaggery extract (20 ml/litre of water) or spray chitin based Pseudomonas fluorescens 20 gram in one litre of water. If the attack is severe, spray Azadirachtin 10 ml or Malathion 2 ml per litre of water.
Coconut	Management	During the beginning of southwest monsoon season, one-kilogram lime can be applied per one matured palm by opening the coconut basin.
Ginger, Turmeric	Planting time	While preparing land for ginger and turmeric cultivations, apply 2 to 3 kg lime per cent. As a prophylactic measure against rhizome rot disease, the ginger and turmeric rhizome should be soaked in the 0.3 percent mancozeb or pseudomonas solution (3 gram per one litter of water) for 30 minutes. Moreover 90 kg FYM mixed with one kg Trichoderma and kept under shade for two weeks with sufficient moisture and can be applied in the basin.
Brinjal	Fruit borer	There is a chance of fruit borer attack in brinjal. It can be controlled by spraying 50 gram neem seed kernel extract per one litre of water. If the attack is severe, spray Chlorantraniliprole @ 3ml per ten litre of water.

Animal Husbandry: - Take precautions to avoid ecto-parasitic infestation. Fever, shivering, brown coloured urine, off feed are noticed, consult veterinary doctor immediately. (CAADECCS, KVASU, Mannuthy).