




Agromet Advisory Service bulletin 17/16, Friday dt.22-04-2016

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara

E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



PALAKKAD

CURRENT SYNOPTIC SITUATION:

An upper air trough runs from Telangana to Comorin area and extends upto 0.9 km above sea level.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 22.04.2016

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS

ENSEMBLE FCST

	DAY-1 23/04	DAY-2 24/04	DAY-3 25/04	DAY-4 26/04	DAY-5 27/04
DISTRICT : PALAKKAD					
Rainfall (mm)	0	0	2	1	0
Max Temperature (deg C)	39	39	39	39	39
Min Temperature (deg C)	27	27	27	27	27
Total cloud cover (octa)	4	4	4	4	4
Max Relative Humidity (%)	80	80	80	80	80
Min Relative Humidity (%)	45	45	45	50	50
Wind speed (kmph)	005	005	004	004	004
Wind direction (deg)	290	290	270	270	250

Agro Meteorological Advisories

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy	Aphid	There is a chance of aphid attack in paddy. Spray Imidacloprid 3 milli in 10 litre of water.
Banana	Sigatoka	Due to the high temperature, there is a chance of sigatoka disease in banana. Spray Pseudomonas 10- 20g per 1 liter of water. In case of severe attack apply SAAF @ 2 g in 1 liter of water.
Coconut		Burying of fresh or dried coconut husk around the palm is a desirable practice particularly for moisture retention. The husk can be buried either in linear trenches taken 3 m away from the trunk between rows of palms or in circular trenches taken around the palm at a distance of 2 m from the trunk. The trenches may be of 0.5 m width and depth. The husks are to be placed in layers with concave surface facing upwards and covered with soil. The beneficial effect of husk burial will last for about 5-7 years. Instead of husk, coconut pith can be buried @ 25 kg / palm / year.