



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



PALAKKAD DISTRICT

AAS 20/2020, (39) Tuesday

12.05.2020

CURRENT SYNOPTIC SITUATION: A cyclonic circulation lies over Comorin area & neighbourhood extending upto 1.5 km above mean sea level. A trough/wind discontinuity runs from West Madhya Pradesh to interior Karnataka across Vidarbha, Marathawada and North Interior Karnataka and extending upto 0.9 km above mean sea level. A cyclonic circulation lies over south Andaman Sea & neighbourhood extending upto mid-tropospheric levels.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	13.05.2020	14.05.2020	15.05.2020	16.05.2020	17.05.2020
Rainfall (mm)	4	6	8	10	5
Maximum temperature(°C)	38	38	38	38	38
Minimum temperature (°C)	27	27	27	27	27
Maximum Relative Humidity (%)	88	88	88	88	88
Minimum Relative Humidity (%)	54	54	54	54	54
Wind speed (kmph)	6	8	10	10	12
Wind direction (deg)	290	270	270	270	270
Total cloud cover (octa)	8	7	8	8	8

As per the forecast of IMD & KAU, there is a chance of moderate rainfall in isolated places of the district. Dried hay, copra, arecanut, etc should be protected from summer rain. Mulching can be practiced to conserve soil moisture.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: As a precaution, use bio-pesticides to control the crop pests and diseases.		
Coconut	Rhinoceros beetle	After thorough cleaning of the crown region, apply 250 g neem cake mixed with equal volume of fine sand (250g) in the inner most 2-3 leaf axils surrounding the spindle leaf. This application has to be repeated thrice a year. (January, May and September/ October) for better results. Or naphthalene balls 12 g (approx. 4 nos) can be put in the innermost 2 leaf axils adjacent to spindle leaf and covered with fine sand, once in 45 days. Or spray <i>Metarrhizium anisopilae</i> (250 ml metarrhizium culture + 750 ml water) in manure pits.
Banana	Pseudo stem weevil	There is a chance of pseudo stem weevil attack in banana. As a precaution, apply 50 g neem cake powder in leaf axils.
Nutmeg	Leaf fall	There is a chance of leaf falling in nutmeg. As a precaution spray 1% bordeaux mixture.
Ginger/Turmeric	-	Prepare land for ginger and turmeric. To avoid foot rot in ginger, sock the suckers in 0.3% mancozeb/ pseudomonas solution for 30 minutes.
Vegetables	-	Addition of pseudomonas culture to the soil at the time of planting of vegetables will help to reduce fungal diseases.