




Agromet Advisory Service bulletin 23/18(46), Friday dt.08.06.2018
(Issued jointly by IAAS Vellanikkara)



Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara
E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

PALAKKAD

CURRENT SYNOPTIC SITUATION: An off shore trough at mean sea level runs from south Maharashtra to North Kerala coast. An East West shear zone runs roughly along Lat 15deg N across south peninsular India between 5.8 & 7.6 km above mean sea level.

A cyclonic circulation lies over East Central Arabian sea off Karnataka Coast between 3.1 & 7.6 km above mean sea level. A cyclonic circulation lies over North Bay of Bengal and neighbourhood between 1.5 & 7.6 km above mean sea level.

Heavy Rainfall warning: Heavy to very heavy rainfall is most likely to occur at isolated places in Kerala till the morning of 9th and isolated heavy rainfall from 9th to the morning of 13th June 2018.

INDIA METEOROLOGICAL DEPARTMENT
NWP MODELS BASED DISTRICT LEVEL WEATHER PREDICTION
ISSUED ON: 08-06-2018
VALID TILL 08:30 IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	WEATHER FCST				
	DAY-1 09/06/2018	DAY-2 10/06/2018	DAY-3 11/06/2018	DAY-4 12/06/2018	DAY-5 13/06/2018
DISTRICT : PALAKKAD					
Rainfall (mm)	45	50	30	30	30
Max Temperature (deg C)	31	31	31	31	31
Min Temperature (deg C)	25	25	25	25	25
Total cloud cover (octa)	7	7	7	7	7
Max Relative Humidity (%)	95	95	95	95	95
Min Relative Humidity (%)	65	65	65	65	65
Wind speed (kmph)	010	010	010	010	010
Wind direction (deg)	250	250	250	250	250

Agro Meteorological Advisories:

Crop	Crop stage / Pest/Disease	Advisories
Paddy	Land preparation to transplanting	Transplant seedlings at 4-5 leaf stage. Soak seed for 30 minutes in a solution of Pseudomonas culture @10g/litre per kg of seeds. The above seed treatment can also be used to control sheath blight, sheath rot, leaf spot disease etc.
Banana	Sigatoka disease	Due to the pre monsoon showers, there is chance of Sigatoka disease in banana. As a precaution, Spray Pseudomonas 20g/litre. In case of severe attack spray 1 gm Bavistin by mixing it with gum in 1 litre of water on lower surface of leaf.
Coconut	Bud rot	Three numbers of perforated Mancozeb sachet (5 gm) can be hung around spindle leaves as a prophylactic measure for bud rot.
Animal Husbandry	Extra care should be taken for wound healing and apply fly repellents around the wound. (CAADECCS, KVASU, Mannuthy)	