



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



**PALAKKAD DISTRICT**

AAS 27/2019(54), Friday

05.07.2019

**CURRENT SYNOPTIC SITUATION:**

A Low Pressure lies over South east Uttar Pradesh & adjoining North east Madhya Pradesh with associated cyclonic circulation extending upto 4.5km above mean sea level. A cyclonic circulation between 3.1 & 7.6 5km above mean sea level lies over North west Bay of Bengal and adjoining Gangetic West Bengal tilting southwards with height.

**Heavy Rainfall Warning:**

**Day 1 : Heavy rainfall** (7- 11 cm in 24 hours) is very likely to occur at isolated places in the district of Kannur .

**Day 2 : Heavy rainfall** (7- 11 cm in 24 hours) is very likely to occur at isolated places in the districts of Kozhikode .

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	06.07.2019	07.07.2019	08.07.2019	09.07.2019	10.07.2019
Rainfall (mm)	20	10	12	15	7
Maximum temperature (°C )	30	30	31	31	31
Minimum temperature (°C )	24	24	25	25	25
Maximum Relative Humidity (%)	98	98	98	98	98
Minimum Relative Humidity (%)	85	85	85	85	85
Wind speed (kmph)	10	10	10	10	10
Wind direction (deg)	270	270	270	270	270
Total cloud cover (octa)	8	7	7	8	6

**Light moderate rainfall is forecasted in the district by IMD & KAU.**

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Pest/ Disease	Advisories
Paddy (Jyothi,Kanchana)	Seedling stage	There is a chance of thrips paddy during the dry spell between monsoon period. Apply 3 ml imidacloprid per 10 litre of water to control them.
Coconut	Bud rot	There is a chance of bud rot in rainy season. Keep three numbers each of perforated Mancozeb sachets (5 gm) around spindle leaves as a prophylactic measure against bud rot.
Pepper	Quick wilt	There is a chance of Quick wilt disease due to high relative humidity. Apply trichoderma or pseudomonas culture in soil or 1% Bordeaux mixture can be sprayed on the leaves and small branches.
Vegetables	Epilachna beetle	There is a chance of Epilachna beetle in vegetables. Cut and remove the leaves along with grub and their eggs. Then apply 2% Neemoil emulsion to control them.
Animal Husbandry	Don't allow the animals to water logged areas to avoid foot lesions. (CAADECCS, KVASU, Mannuthy)	

