


Agromet Advisory Service bulletin 34/16, Tuesday dt 16-08-16

(Issued jointly by IAAS Vellanikkara)





PALAKKAD

CURRENT SYNOPTIC SITUATION: A well marked low pressure area lies over northwest Bay of Bengal and neighbourhood. Associated upper air cyclonic circulation extends upto 5.8 km asl. A feeble trough of low pressure lies over Lakshadweep area.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 16.08.2016 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 17/08	DAY-2 18/08	DAY-3 19/08	DAY-4 20/08	DAY-5 21/08
DISTRICT : PALAKKAD					
Rainfall (mm)	6	9	9	6	5
Max Temperature (deg C)	30	30	30	30	30
Min Temperature (deg C)	23	23	23	23	23
Total cloud cover (octa)	6	6	6	6	6
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	008	800	008	008	008
Wind direction (deg)	290	290	270	270	270

Agro Meteorological Advisories

G	Stage of the			
Crop	crop /	Advisories		
	Pest/Disease			
		There is a chance for Sheath blight disease during rainy season.		
Paddy	Panicle	The immediate symptom is that outer leaves of paddy become		
(Jyothi,	initiation/	yellow coloured. Black lesions can be seen on the lower part		
Kanchana)	Sheath blight	just above the water level. Apply Propiconazole 25 EC @ 500		
		ml per 1 hector to control Sheath blight.		
		There is a chance of yellowing in banana. Examine the sucker		
Banana	Yellowing	and make sure that it is Erwinia rot, it can be controlled by		
	(Erwinia Rot)	mixing 25gms of Kocide and 6gms of Streptocycline in 25 ml		
		of water and apply 5 litre of solution to each banana basin.		
		During rainy season there is a chance of rhizome rot, to control		
Ginger, Turmeric	Rhizome rot	this dig out the affected plants and apply Kocide 2gm/1 litre		
		water.		
		To control fruit borer Apply any pesticide contains Neem oil (4		
Cowpea	Fruit borer	ml per one liter of water).Repeat it in every week.		