



Advisory Service bulletin 35/16, Friday dt.26-08-2016

(Issued jointly by IAAS Vellanikkara)





PALAKKAD

CURRENT SYNOPTIC SITUATION: A low pressure area has formed over west central Bay of Bengal off North Andhrapradesh coast. A trough of low pressure on sea level chart extends from Karnataka coast to Kerala coast.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 26.08.2016 VALUE THE 1.08.20 HDS IST OF THE NEXT 5 DAYS

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 27/08	DAY-2 28/08	DAY-3 29/08	DAY-4 30/08	DAY-5 31/08
DISTRICT : PALAKKAD					
Rainfall (mm)	14	6	6	7	8
fax Temperature (deg C)	32	32	32	32	31
fin Temperature (deg C)	24	24	24	24	24
otal cloud cover (octa)	7	7	7	7	7
Max Relative Humidity (%)	90	90	90	90	90
in Relative Humidity (%)	70	70	70	70	70
ind speed (kmph)	008	800	800	006	006
Ind direction (deg)	290	290	290	290	270

Agro Meteorological Advisories

Crop	Stage of the crop / Pest/Disease	Advisories	
D 11	D : 1 /	Apply neem based pesticides (3-4 ml per one liter of	
Paddy	Panicle stage/	water). Pesticides application should be done either before	
(Jyothi, kanjana)	Rice bug	9 am or after 3 pm. Otherwise, the pesticide application	
		will adversely affect the pollination.	
		There is a chance of bud rot disease during rainy season.	
Coconut	Bud rot	Cut and remove all the affected tissues of the crown.	
	disease	Apply Bordeaux paste and protect it from rain till normal	
		shoot emergence.	
Banana	Sigatoka	Due to the changing weather pattern, there is a chance of sigatoka disease in banana. As a precaution, Spray Pseudomonas 20g per 1 liter of water. In case of severattack spray 1 gm Bavistin by mixing it with gum in 1 litro of water on lower surface of leaf.	
Cowpea	Bug Apply neem based pesticides 3-4 ml per one liter of w Spray malathion 2 ml/ 1 litre of water.		