





Agromet Advisory Bulletin for PALAKKAD District (Valid from 19.10.2024 to 23.10.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 42/2024, (84) Friday

Dated 18.10.2024

A. Medium range weather forecast of IMD for next five days

Weather parameters	19.10.2024	20.10.2024	21.10.2024	22.10.2024	23.10.2024
Rainfall (mm)	2	1	0.2	0.2	0.2
Maximum temperature (°C)	32	32	32	32	32
Minimum temperature (°C)	25	25	24	24	24
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	4	5	4	3	4
Wind direction (deg)	290	290	250	320	320
Total cloud cover (octa)	3	4	3	3	3
Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in					
Weather summary / Alert: A	-			nance of ligh	t rai

the district. Below normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agro meteorological Advisories

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports to banana crops. Make the vegetable pandals stronger.

0 1	0		
Сгор	Pest/ Disease, Crop Management	Agromet Advisory	
Rice (Mundakan) Seedling stage	Transplanting	As a prophylactic measure, the root of seedlings can be soaked for 30 minutes in a solution of pseudomonas culture to control sheath blight, sheath rot, leaf spot disease etc. Spray 20 gram Pseudomonas in one litre of water before transplanting.	
Ginger	Rhizome rot	There is a chance of rhizome rot disease in ginger. As a prophylactic measure spray 2 gram Copper oxychloride or 1.5 gram copper hydroxide per one litre of water.	
Cucurbitaceous Vegetables	Downy mildew	During high relative humidity, there is a chance of downy mildew attack in cucurbitaceous vegetables. Spray 2.5gm Mancozeb per one litre of water under the leaves.	
Animal Husbandry: - Cattle feed should be stored on a wooden board or iron board. If moisture			

Animal Husbandry: - Cattle feed should be stored on a wooden board or fron board. If moisture is in the cattle feed it may cause deadly fungal growth. Vaccinate animals against foot and mouth disease. (CAADECCS, KVASU, Mannuthy).