



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



PALAKKAD DISTRICT

AAS 45/2018(89), Friday

09.11.2018

The Agricultural advisory of this bulletin is not applicable in flood-affected areas.

CURRENT SYNOPTIC SITUATION: The low pressure over Andaman sea and adjoining Malay Peninsula now lies over central parts of Andaman sea and neighbourhood with associated cyclonic circulation extending upto mid Tropospheric levels. The cyclonic circulation over Comorin area and adjoining Sri Lanka and Equatorial Indian ocean extends upto 3.1 km above mean sea level .

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

| Weather parameters | 10.11.2018 | 11.11.2018 | 12.11.2018 | 13.11.2018 | 14.11.2018 |
|-------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Maximum temperature (°C) | 33.1 | 33.2 | 33.2 | 33.2 | 33.2 |
| Minimum temperature (°C) | 24.2 | 24.3 | 24.3 | 24.3 | 24.3 |
| Total cloud cover (octa) | 5 | 4 | 4 | 4 | 3 |
| Maximum Relative Humidity (%) | 80 | 80 | 80 | 80 | 80 |
| Minimum Relative Humidity (%) | 53 | 41 | 43 | 45 | 45 |
| Wind speed (kmph) | 10 | 10 | 10 | 10 | 10 |
| Wind direction (deg) | 70 | 70 | 70 | 70 | 70 |

As per IMD forecast, there is less chance for rainfall for the coming days in the district.

AGRO-METEOROLOGICAL ADVISORIES:

| Crop | Crop stage | Pest/ Disease | Advisories |
|-----------------------|---|----------------------------|--|
| Pepper | - | Yellowing | Yellowing was observed in pepper due to flood situation followed by physiological drought. Soil drenching with 2g kocide per one litre of water, followed by a foliar spray and soil drenching with 20 g of pseudomonas per litre of water, after two weeks was recommended. |
| Banana and Vegetables | - | Spodoptera | Destroy the affected leaves along with the spodoptera. If spodoptera attack is severe, spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml per 10 litre of water. |
| Paddy | Seedling stage | Leaf folder and Stem borer | As a prophylactic measure, apply 2CC <i>Trichogramma chilonis</i> card per acre to control leaf folder and 2CC <i>Trichogramma japonicum</i> card per acre to control stem borer in paddy. Cut the cards in to small pieces and fix in different parts of the field in plastic cups. If the attack is severe, spray 3ml chlorantraniliprole per 10 litre of water. |
| Animal Husbandry | Give FMD vaccination to cattle and goats and Leptospira vaccination to pet animals. (CAADECCS, KVASU, Mannuthy) | | |