



Agromet Advisory Service bulletin 49/17, Friday dt. 08-12-2017

(Issued jointly by IAAS Vellanikkara)





PALAKKAD

CURRENT SYNOPTIC SITUATION:

The depression over South east Bay of Bengal and neighbourhood intensified into a Deep Depression over central Bay of Bengal moved north northwestwards and lies over west central Bay of Bengal near lat 15° N and long 6.8° E.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 08.12.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

DISTRICT : PALAKKAD ENSEMBLE FCST

PARAMETERS	09.12.2017	10.12.2017	11.12.2017	12.12.2017	13.12.2017
Rainfall (mm)	3	0	0	0	0
Max Temperature (deg	34	34	34	34	34
Min Temperature (deg C)	23	23	23	23	23
Total cloud cover (octa)	4	4	4	4	4
Max Relative Humidity (%)	85	85	85	85	85
Min Relative Humidity (%)	55	55	55	55	55
Wind speed (kmph)	10	10	10	10	10
Wind direction (deg)	90	90	90	90	90

Agro Meteorological Advisories:

Стор	Crop stage / Pest/Disease	Advisories		
Paddy (Jyothi, Kanchana)	Rice bug	Apply neem based pesticides (3ml per one liter of water). Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.		
Coconut	Bud rot	As a prophylactic measure, hung three perforated packets of Mancozeb sachet (5gm) around spindle leaves.		
Banana	Spodoptera	Destroy the affected leafs along with the spodoptra. For severe attack spray Flubendiamide (Fame) 2 ml/10 L of water.		
Vegetables (Snake gourd, Bitter gourd)	Fruit fly	Remove and destroy affected and decayed fruits. Use special pheromone traps like cue lure (6 nos/Acre) for vegetable fruit fly or Thulasi/fruit traps. If the attack is not reduced, spray mixture of malathion 2 ml per one litre of water + jaggery 10 g at flowering and fruit initiation.		
Animal Husbandry	If respiratory distress, fever or lameness are noticed in animals, consult a veterinary surgeon immediately. (CAADECS, KVASU, Mannuthy)			