

Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin





PALAKKAD DISTRICT

AAS 51/2018(101), Friday

21.12.2018

CURRENT SYNOPTIC SITUATION: The trough of low pressure at mean sea level over central parts of south Bay of Bengal now lies over southwest Bay of Bengal and adjoining equatorial Indian Ocean and extends upto 2.1km above mean sea level. The cyclonic circulation over southeast Arabian Sea and adjoining Lakshadweep area has become less marked.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

| Weather parameters | 22.12.2018 | 23.12.2018 | 24.12.2018 | 25.12.2018 | 26.12.2018 |
|-------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 0 | 7 | 0 | 0 | 0 |
| Maximum temperature (°C) | 32.1 | 32.1 | 32.1 | 32.1 | 32.1 |
| Minimum temperature (°C) | 24.1 | 24.1 | 24.1 | 24.1 | 24.1 |
| Total cloud cover (octa) | 4 | 6 | 4 | 4 | 4 |
| Maximum Relative Humidity (%) | 85 | 85 | 85 | 85 | 85 |
| Minimum Relative Humidity (%) | 45 | 45 | 45 | 45 | 45 |
| Wind speed (kmph) | 12 | 14 | 12 | 8 | 12 |
| Wind direction (deg) | 90 | 90 | 90 | 110 | 270 |

No rainfall is forecasted for coming five days in the district by India Meteorological Department.

AGRO-METEOROLOGICAL ADVISORIES:

| Crop | Crop stage | Pest/ Disease | Advisories | | |
|---------------------|---|---------------|--|--|--|
| Paddy | Tillering stage | Leaf folder | As a prophylactic measure, apply 2CC <i>Trichogramma chilonis</i> card per acre to control leaf folder. Cut the cards in to small pieces and fix in different parts of the field in plastic cups. | | |
| Vegetables (cowpea) | Fruiting stage | Fusarium wilt | To prevent Fusarium wilt in cowpea, mix 20 g of cow dung in one litre of water and soil should be drenched with the supernatant liquid by mixing with 20 g of pseudomonas. | | |
| Banana | - | Spodoptera | Destroy the affected leafs along with the <i>Spodoptera</i> . If <i>Spodoptera</i> attack is severe, spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml per ten litre of water. | | |
| Animal Husbandry | Provide plenty of clean drinking water in the cattle sheds and grazing lands. (CAADECCS, KVASU, Mannuthy) | | | | |