



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



PALAKKAD DISTRICT

AAS 52/2018(103), Friday

28.12.2018

CURRENT SYNOPTIC SITUATION: A cyclonic circulation extending upto 0.9km above mean sea level lies over Comorin area and neighbourhood.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	29.12.2018	30.12.2018	31.12.2018	01.01.2019	02.01.2019
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	34	34	34	34	34
Minimum temperature (°C)	23	23	23	23	23
Total cloud cover (octa)	6	5	4	4	4
Maximum Relative Humidity (%)	85	85	85	85	85
Minimum Relative Humidity (%)	40	40	40	40	40
Wind speed (kmph)	6	6	6	6	6
Wind direction (deg)	110	110	110	110	110

No rainfall is forecasted for coming five days in the district by India Meteorological Department.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Crop stage	Pest/ Disease	Advisories
Paddy	Tillering stage	Leaf folder	As a prophylactic measure, apply 2CC <i>Trichogramma chilonis</i> card per acre to control leaf folder. Cut the cards in to small pieces and fix in different parts of the field in plastic cups.
Cucurbitaceous Vegetables	-	Powdery mildew	White powdery spots are seen on the leaves of the affected plants. If any fruits are present, harvest them. Then apply 1.5 ml tebuconazole or 0.5 ml difenoconazole per one litre of water. After 10 days, apply 20 g of <i>Trichoderma viride</i> per one litre of water to control powdery mildew disease of vegetables.
Banana	-	<i>Spodoptera</i>	Destroy the affected leafs along with the <i>Spodoptera</i> . If <i>Spodoptera</i> attack is severe, spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml per ten litre of water.
Animal Husbandry	Provide plenty of clean drinking water in the cattle sheds and grazing lands. (CAADECCS, KVASU, Mannuthy)		