

## Kerala Agricultural University District Agrometeorological Unit Krishi Vigyan Kendra, Palakkad

Phone: 0466 2212279 e-mail: kvkpalakkad@kau.inAgromet



**Advisory Service Bulletin** 

## **Palakkad**

Issued jointly by Kerala Agricultural University, India Meteorological Department, Palakkad District Agrometeorological Unit and Department of Agriculture

AAS. 99/2022 (Tuesday)				Date: 13.12.2022			
WEATHER FORECAST FOR NEXT FIVE DAYS							
Weather parameters	14.12.2022	15.12.2022	16.12.2022	17.12.2022	18.12.2022		
Rainfall (mm)	6	5	0	0	0		
Maximum temperature (°C)	29	30	32	32	32		
Minimum temperature (°C)	23	22	22	22	22		
<b>Maximum Relative Humidity (%)</b>	95	95	95	95	95		
9Minimum Relative Humidity (%)	75	75	75	75	75		
Wind speed (km/hr))	2	3	3	3	4		
Wind direction (degree)	340	90	70	70	70		
Cloud cover (okta)	8	8	5	3	3		

Weather summary / Alert: Light rainfall is expected in the district. Normal rainfall is expected in Kerala.

General Advisory: Prophylactic measures to control crop pest & diseases can be taken to avoid their spread. Proper mulching can be followed to conserve soil moisture. e.

## **Advisory**

Crops	Pests/ diseases	Crop advisory				
Rice (Mundakan) Tillering stage Bacterial	Leaf folder	During cloudy weather condition, there is a chance of Leaf folder attack in Mundakan Paddy. It can be controlled by using trichogramma cards (2CC per one acre). Cut the cards in to small pieces and fix in different regions of the field. If the attack is severe, spray Chlorantraniliprole @ 3ml per ten litre of water.				
		As a prophylactic measure to manage Bacterial Leaf Blight (B L B), dissolve 20 g of cow dung in one litre of water and spray the supernatant liquid by mixing with 20g of pseudomonas. Apply bleaching powder @ 20 bags of 100 gm each for one acre in the irrigation channels for checking the spread of the disease. Spry antibiotics viz. 6 g of streptocycline in 20 litre of water in severe cases.				
Vegetables (Cowpea)	Anthracnose disease	There is a chance of Anthracnose disease in Cowpea. As a prophylactic measure, apply 1% Bordeaux mixture or spray pseudomonas 20 gm pe liter. Pesticides should not be applied during rain				
Cucurbitaceous Vegetables	Downy mildew	Due to high relative humidity, there is a chance of downy mildew attack in cucurbitaceous vegetables. Spray 2.5gm Mancozeb per one litre of water under the leaves. Pesticides should not be applied during rain.				

Animal Husbandry- Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospirosis Fever, shivering, and brown coloured urine or any other symptoms of jaundice if noted, immediately consult a veterinary doctor.