



**Kerala Agricultural University, Vellanikkara, Thrissur**  
**Department of Agricultural Meteorology, College of Horticulture**

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

**Agromet Advisory Service Bulletin**

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



**THRISSUR DISTRICT**

**AAS 02/2020 (04), Friday**

**10.01.2020**

**SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:**

The total rainfall received 0.0 mm. The maximum temperature ranges from 34.1°C to 34.9°C, minimum temperature ranges from 22.6°C to 24.2°C. Morning relative humidity ranges from 76 % to 95% and afternoon relative humidity ranges from 41% to 50%. Wind speed ranges from 1.8 to 9.6 km/h. Bright sunshine hours range from 9.0 to 9.9 hours. Evaporation ranges from 3.4 to 5.9 mm.

**CURRENT SYNOPTIC SITUATION:** Northeast monsoon rains have ceased over Tamil Nadu, Puducherry& Karaikkal, Kerala & Mahe & adjoining areas of coastal Andhra Pradesh & Yanam, Rayalaseema and south Interior Karnataka from today the 10<sup>th</sup> Jan 2020. Relatively dry air prevails over Kerala.

**WEATHER FORECAST FOR THE NEXT FIVE DAYS:**

Weather parameters	11.01.2020	12.01.2020	13.01.2020	14.01.2020	15.01.2020
Rainfall (mm)	0	0	0	0	0
Maximum temperature(°C )	33	33	33	33	33
Minimum temperature (°C )	21	21	21	21	21
Maximum Relative Humidity (%)	72	72	72	72	72
Minimum Relative Humidity (%)	47	47	47	47	47
Wind speed (kmph)	12	16	10	10	10
Wind direction (deg)	50	50	50	50	50
Total cloud cover (octa)	3	2	1	4	3

**As per the forecast of IMD & KAU, due to no rainfall and prevailing dry weather conditions, irrigation of crops to be continued in the district.**

**AGRO-METEOROLOGICAL ADVISORIES:**

Crop	Pest/ Disease	Advisories
<b>General Advisories:</b> As a precaution, use bio-pesticides to control the crop pests and diseases.		
Rice (Mundakan) Panicle Emergence	Stem borer	As a prophylactic measure, apply 2CC Trichogramma japonicum card per acre to control stem borer in paddy. Cut the cards in to small pieces and fix in different parts of the field. If the attack is severe, spray 3ml chlorantraniliprole per 10 litre of water.
	Bacterial Leaf blight	To control the spread of bacterial leaf blight in paddy, spray 6g of streptocycline in 30 litre of water.
Vegetables (cowpea)	Fusarium wilt	There is a chance of fusarium wilt in cowpea. Drench the soil rhizosphere with 2g of bavistin or 2 ml of contaf per one litre of water alternatively within 15 days interval to control the disease. Severely affected plants should be removed and destroyed from the field and apply a pinch of bleaching powder in the pit.
Chilli	White fly	There is a chance of white fly attack in chilli. Apply 2% neem oil garlic emulsion or spray 20g <i>Lecanicillium lecanii</i> per one litre of water in ten days intervals.
Animal Husbandry		Vaccinate animals against foot and mouth disease and hemorrhagic septicemia. (CAADECCS, KVASU, Mannuthy)