

Agromet Advisory Service bulletin 05/16 Tuesday dt 26-01-2016

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara


E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

THRISSUR

Significant weather for the preceding week at Vellanikkara



Rainfall weekly total	23.8 mm
Maximum temperature	30.2-34.3 ⁰ C
Minimum temperature	22.0-25.6 ⁰ C
Max. RH	63-96%
Min. RH	40-63%
Wind speed	1.2-8.8 km/hr
Sun shine hour	0.0-9.1 hr
Evaporation	1.6-6.0 mm



CURRENT SYNOPTIC SITUATION: An upper air cyclonic circulation lies over Maldives and Lakshadweep area extending upto 0.9 km above sea level.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 25.01.2016

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS

ENSEMBLE FCST

	DAY-1 26/01	DAY-2 27/01	DAY-3 28/01	DAY-4 29/01	DAY-5 30/01
Rainfall (mm)	0	2	0	0	0
Max Temperature (deg C)	34	34	34	34	34
Min Temperature (deg C)	24	24	24	24	24
Total cloud cover (octa)	5	5	3	3	2
Max Relative Humidity (%)	80	80	80	75	75
Min Relative Humidity (%)	50	50	45	40	40
Wind speed (kmph)	006	006	008	008	008
Wind direction (deg)	50	50	50	50	50

DISTRICT : THRISSUR

Crop stage

This is the time for the preparation of punja field. Banana plant is 3 month after planting. This is the time for starting irrigation in coconut and arecanut. It is the time for starting mulching in coconut and pepper.

Agro Meteorological Advisories

Crop	Pest/Disease	Advisories
Paddy	Land preparation to transplanting	Plough the land for Punja cultivation. Transplant seedlings at 4-5 leaf stage. Soak seed for 30 minutes in a solution of Pseudomonas culture @10g/litre of water per kg of seed. The above seed treatment can also be used to control sheath blight, sheath root, leaf spot disease etc.
Coconut	Mite	Due to high temperature, there is a chance of increase of mite population in coconut. Apply 2% neem oil-garlic-soap emulsion in the crown on young bunches to prevent mite's attack.
Banana	Sigatoka	Spray Pseudomonas at the rate of 15g per one liter of water or SAAF at the rate of 2 g in one liter of water.