



Gramin Krishi Mousam Sewa (GKMS), Thrissur



Agromet Advisory Bulletin for Thrissur District

(Valid from 26.2.2022 to 2.3.2022)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 08/2022, (16) Friday

dated 25.2.2022

A. Significant weather for the preceding week at vellanikkara								
Rainfall	Maximum	Minimum	RH	RH	Wind speed	Sun shine	Evap. (mm)	
(mm)	Temp.(°C)	Temp.(°C)	FN (%)	AN (%)	(Km/h)	hours		
(0.0)	(34.3-36.1)	(22.2-24.6)	(66-93)	(30-55)	(2.0-8.0)	(5.7-9.9)	(3.4-7.1)	

B. Medium range weather forecast of IMD for next five days						
Weather parameters	26.2.2022	27.2.2022	28.2.2022	01.3.2022	02.3.2022	
Rainfall (mm)	0	0	0	0	0	
Maximum temperature (°C)	36	36	36	36	36	
Minimum temperature (°C)	24	24	24	24	24	
Maximum Relative Humidity (%)	70	70	70	70	70	
Minimum Relative Humidity (%)	30	30	30	30	30	
Wind speed (kmph)	2	3	2	2	2	
Wind direction (deg)	270	270	270	270	270	
Total cloud cover (octa)	2	3	3	7	6	

Weather summary / Alert: Since dry weather condition is prevailed, irrigation of crops can be continued, when required in the district. Below normal rainfall is expected in Kerala.

RAINFALL FORECAST FOR NEXT FIVE DAYS							
2022 February 26	2022 February 27	2022 February 28	2022 March 01	2022 March 02			
No Rain							

C. General and Agrometeorological Advisories for Thrissur District Since dry weather condition is prevailed, irrigation of crops can be continued. Pest/ Disease Crop Crop/ Animal Advisory To control Blast disease in rice, spray 4 gram of Nativo Puncha Rice (Trifloxystrobin + Tebuconazole) mixture in ten litres of in Active **Blast** water. Tillering stage There is a chance of white fly attack in Brinjal. Apply 2% White fly neem oil garlic emulsion or spray 20g Lecanicillium lecanii **Brinjal** per one liter of water in ten days intervals. There is a chance of thrips attack in chilli. Apply 2% neem oil+ garlic emulsion under the leaf surface at 10 days **Thrips** intervals. If the attack is severe, apply spiromecifen 8 ml per Chilli ten litre of water. Ensure sufficient supply of clean drinking water to the animals. Take measures for Animal Husbandry preventing mosquito breeding in the premises. (CAADECCS, KVASU, Mannuthy).